

Final Report - NCCP/MSCP Raptor Monitoring
Project (January 1, 2001 – December 31, 2003)

for

**California Department Fish and Game
4949 Viewridge Ave.
San Diego, CA 92123**

by

**Wildlife Research Institute, Inc.
P.O. Box 2209
18030 Highland Valley Road
Ramona, CA 92065**

March 31, 2005

TABLE OF CONTENTS

BACKGROUND	1
THE PROJECT AND ITS OBJECTIVES	1
THIS REPORT	2
METHODS	2
RESULTS	7
DISCUSSION	13
RECOMMENDATIONS	19
ACKNOWLEDGEMENTS	25
LITERATURE CITED	26
APPENDIX A - Project Bibliography	
APPENDIX B - Breeding Season Raptor Nests and Territories by Site (2001-2003)	
APPENDIX C - Proposed Long-term Raptor Monitoring Program	

LIST OF TABLES AND FIGURES

Figure 1 – MSCP Raptor Study Site Locations	4
Figure 2 – Proposed MSCP Raptor Monitoring Areas (Breeding Season)	22
Figure 3 – Proposed MSCP Winter Raptor Monitoring Area	25
Table 1. MSCP Study Sites Chosen to be Surveyed for Raptors	5
Table 2. Number of Raptor Nests and/or Territories by Site (2001-2003)	9
Table 3. Number of Raptors Observed During the Winter Surveys (2001-2003)	10
Table 4. Management and Enforcement Issues by Raptor Survey Site	15
Table 5. Proposed MSCP Areas for Long-term Raptor Monitoring (Breeding Season)	21

BACKGROUND

The Natural Communities Conservation Planning (NCCP) Raptor Monitoring Project is part of the urgent implementation tasks associated with the Multiple Species Conservation Program (MSCP). The MSCP is the local representation of the State's NCCP Program of which the City of San Diego is a participating member and the lead agency. The County of San Diego is also an active participant (County of San Diego 1997). The city adopted the MSCP on March 18, 1997 and entered into a binding contract on July 16, 1997 with the State of California Department of Fish and Game and the United States Fish and Wildlife Service to implement the MSCP.

Each habitat conservation plan (HCP) requires a monitoring program to determine the efficacy of that plan. The "Biological Monitoring Plan for the Multiple Species Conservation Program" (Ogden 1996) recommended monitoring for certain plant species, coastal sage scrub (Coastal California Gnatcatcher and Cactus Wren), herpetofauna, and grasslands (specifically, using raptors).

THE PROJECT AND ITS OBJECTIVES

Monitoring of raptors is a critical component of the MSCP. This project, specifically, addresses monitoring the raptor species identified as target species for MSCP monitoring with one exception--the Burrowing Owl (BO; *Athene cunicularia hypugaea*). In addition to the Burrowing Owl, the MSCP Biological Monitoring Plan (Ogden, 1996) identified the following raptor species (hereafter referred to as the "target" species) to be monitored: Golden Eagle (GE; *Aquila chrysaetos*), Bald Eagle (BE; *Haliaeetus leucocephalus*), Peregrine Falcon (PF; *Falco peregrinus*), Northern Harrier (NH; *Circus cyaneus*), Ferruginous Hawk (FH; *Buteo regalis*), Swainson's Hawk (SH; *Buteo swainsoni*), and Cooper's Hawk (CH; *Accipiter cooperii*). Prior to the subject work, no comprehensive study had been conducted for any of these species, within the geographical limits of the MSCP.

The Wildlife Research Institute, Inc. (WRI), a non-profit organization, has been working with all MSCP participants to identify appropriate long-term raptor monitoring locations (based on the results of the current WRI raptor surveys), develop a scientifically-based monitoring program (including survey locations and protocols), test the monitoring methods, and identify opportunities for population enhancements.

The original project objectives (taken from the contract's scope of work) are as follows:

- Determine where breeding and wintering individuals (of the target species) are located within the study areas.
- Wherever possible, document the breeding success of active pairs.
- Characterize situations of both successful and less successful or unsuccessful habitat.
- Identify, modify, or create, if necessary, survey raptor monitoring methods, based on scientific principles that would be appropriate to meet the objectives of the MSCP Monitoring Plan.
- Identify management, including research, needs and enhancement opportunities.

THIS REPORT

Constraints. This report covers WRI's raptor surveying activity for the three years of this project (January 1, 2001 through December 31, 2003), focusing on the breeding and wintering seasons. For the record, our work did not, officially, include the BO. Therefore, with few exceptions, surveys were not conducted during what would normally have been the most productive time for this species (i.e., early morning and early evening). Fieldwork was conducted during the daylight hours to maximize chances for seeing the diurnal raptors that were the focus of the contracted scope. Although nocturnal owls can be expected to nest and winter in many of the study sites, they would be expected to often escape observation under this temporal survey regime. However, our methods required documenting any raptor, regardless of whether or not it was a target species and, when a BO or any other owl was observed, it was noted.

A natural phenomenon created a situation that could be considered a constraint. This was the extreme drought that the region experienced for several years (1999-2004). Therefore, 2001 through 2003 may not have been the best of raptor breeding years. Drought clearly plays a significant factor in the density and reproductive success of raptors. This study was conducted during the worst drought for San Diego in over 160 years. This fact should be noted for future researchers and resource managers/planners. This kind of extreme drought has the potential effect of reducing the available prey biomass, which, in turn, can have at least two effects. First, it likely reduces the "attractiveness" of a habitat complex, partly because of low prey densities, and may encourage raptors and other predators to look elsewhere. Second, for those individuals that choose to stay in a less-than-ideal environment, the lack of prey often results in lowered reproductive success or even total nest failure (see Discussion, below). If a nest site is not successful, the birds are more likely to disperse, which leaves the historically active territories apparently, or actually, vacant.

Intent. It is the intent that this, the Final Report, will not only serve to (1) provide data analysis and interpretation but, importantly, it strives to (2) provide an initial baseline of information on many of the breeding and wintering raptors within the MSCP and environs, (3) identify resource management challenges and opportunities, and (4) recommend needed research and management, including what areas should be considered for the MSCP Long-term Raptor Monitoring Program (LRMP).

METHODS

LITERATURE REVIEW, INTERVIEWS, DATA SEARCHES, ETC.

We first contacted other professional biologists, regarding available literature and monitoring programs already in place. We acquired relevant literature, which we did not already have, and met with and/or phone-interviewed members of the outdoor-oriented public as well as key professionals in the San Diego ornithologist community (including Mr. John Oakley, Mr. David Mayer, Mr. Phil Unitt, Dr. Jim Hannan, and others listed in the Acknowledgements section) to inquire about raptor sightings. Using existing published and gray literature, the Natural Communities Data Base, museum collections, raw data from the San Diego County Bird Atlas (then in prep.), MSCP vegetation and sensitive species GIS data, and discussions with knowledgeable experts, a project bibliography, relevant to the MSCP and the target species, was produced (Appendix A).

STUDY SITES

The choice of *study sites* (i.e., those which would be the focus of the 2001-2003 field observations) began with the raptor monitoring locations proposed by the “Biological Monitoring Plan for the Multiple Species Conservation Program” (Ogden 1996). Through consultations with CDFG staff and other knowledgeable biologists, we initially identified 22 sites. After some consolidation and the addition of several sites, including control sites and five sites recently acquired by the state or federal government (numbers 34, and 39 through 43), this number was, ultimately, increased to 45 locations within, and juxtaposed to, the MSCP (hereafter referred to as “study sites”; Figure 1 and Table 1). These became the sites, which were surveyed and considered as *potential* sites, or components of sites, for the Long-term *Monitoring Plan*. The basis for choosing the study sites included that they (1) could be expected to support raptors, (2) were part of an area which was managed by a public or private organization or, alternatively, could serve as a control site over time, (3) were accessible by vehicle and could be safely surveyed with repeatability, (4) contained grassland and/or other relevant habitat which was representative of the MSCP area, and (5) were within or immediately juxtaposed to the MSCP area. We considered all ten sites recommended by the Ogden (1996) report. Of those ten sites, we believe all are covered by one or more of the above 45 locations unless they did not meet the above criteria.

MONITORING SITES

The parameters considered in order to make the recommendations for monitoring sites (i.e., those which would be used in the MSCP Long-term Monitoring Program; LRMP) were discussed at a meeting with representatives of CDFG, USFWS, the City of San Diego and the County of San Diego, on January 27, 2002, at the CDFG San Diego office. It was agreed that the following were important when reviewing each study site as a potential MSCP LRMP site:

- Number of individual raptors documented at a site
- Number of raptor species
- Number of target raptor species
- Diversity of raptors and/or target raptor species
- Number of raptor territories
- Number of crows and/or ravens
- Incidence and/or expectation of management/enforcement problems
- Likely changes in habitat and disturbance over time

In order to identify which sites are the most appropriate for the MSCP LRMP during the breeding season, each site was examined, based on two species diversity parameters (number of total raptors and number of target raptors, both of which were normalized by level of effort) and a third parameter for evenness (Probability of an Interspecific Encounter or PIE; Hurlburt, 1971). The analysis for evenness provided a logical break between the top 19th and 20th sites. All sites were then arranged in descending order for each of these three parameters. If any site came out in the top 19 for any two of the three parameters, it was considered a candidate for the MSCP LRMP. Seventeen sites met this requirement. Each site was reviewed, based on our biological knowledge of that site and how it fit into the geographic distribution of recommended monitoring sites. Finally, juxtaposed sites were combined and sites and site boundaries were adjusted based on historic raptor numbers and improved geographic coverage.




Table 1. Raptor Study Sites (2001-2003)

NOTE TO READER: In order to facilitate the reader's access to the following topographic maps, they are listed below alphabetically and by site number.

<u>Number</u>	<u>Name</u>	<u>Name</u>	<u>Number</u>
1	Crestridge	Boden Canyon	2
2	Boden Canyon	Border Fields	44
3	Jamul Ranch	Brown Field Complex	22
4	SDNWR*/Salt Works/Egger Ghio	Crestridge	1
5	McGinty Mountain Complex	Grasslands/Route 67	30
6	San Diego Bay NWR (winter only)	Hollenbeck Canyon	34
7	Lake Hodges	Immenschuh	39
8	Penasquitos Lagoon	Iron Mountain	11
9	Torrey Pines	Jamul Ranch	3
10	Sycamore Canyon	Lake Hodges	7
11	Iron Mountain	Los Montanas (North)	40
12	Otay Mountain	Los Montanas (South)	41
13	Marron Valley	Marron Valley	13
14	Otay Lakes	McGinty Mountain Complex	5
15	SDNWR* Sweetwater Marsh	Miramar Reservoir	20
16	San Vicente	Mission Bay	21
17	Sycuan Peak	Mission Trails	24
18	Point Loma	North Island	19
19	North Island	Otay Lakes	14
20	Miramar Reservoir	Otay Mountain	12
21	Mission Bay	Penasquitos Canyon	33
22	Brown Field Complex	Penasquitos Lagoon	8
23	SDNWR*/San Miguel Mountain	Point Loma	18
24	Mission Trails	Proctor Valley	25
25	Proctor Valley	Rancho San Diego (East)	42
26	San Diego River	Rancho San Diego (West)	43
27	Route 67 South	Rock Mountain	35
28	San Dieguito Lagoon	Rockwood Canyon	32
29	Route S-6 (deleted/safety issue)	Route 67 South	27
30	Grasslands/Route 67	Route 94 (North and South)	38
31	Sloan Canyon	Route S-6	29
32	Rockwood Canyon	San Diego Bay NWR (winter only)	6
33	Penasquitos Canyon	San Diego River	26
34	Hollenbeck Canyon	San Dieguito Lagoon	28
35	Rock Mountain	San Pasqual	36
36	San Pasqual	San Vicente	16
37	SDNWR*Tijuana Slough	SDNWR* Sweetwater Marsh	15
38	Route 94 (North and South)	SDNWR*/Salt Works/Egger Ghio	4
39	Immenschuh	SDNWR*/San Miguel Mountain	23
40	Los Montanas (North)	SDNWR*Tijuana Slough	37
41	Los Montanas (South)	Sloan Canyon	31
42	Rancho San Diego (East)	Sweetwater Reservoir	45
43	Rancho San Diego (West)	Sycamore Canyon	10
44	Border Fields	Sycuan Peak	17
45	Sweetwater Reservoir	Torrey Pines	9
*San Diego National Wildlife Refuge			

After completing the above analysis, it became clear that the coastal portions of the MSCP were excluded from the proposed breeding season monitoring because the vast majority and greatest diversity of raptor species breed somewhat inland of the coast. In addition, our data showed that the MSCP area supported a sizable wintering PF population, most of which would be excluded without a coastal component to the MSCP LRMP. Therefore, a winter monitoring route was established that included a good sampling of the coastal wintering raptor habitat that could be driven safely and consistently.

FIELD SURVEYS

By way of clarification, we will be discussing two kinds of raptor searching and documentation. The first is the *survey*—the approach we took to investigate each of the 45 study sites, some of which we are recommending for the MSCP LRMP. This approach utilized several techniques in order to capture a maximum amount of raptor data on sites of considerable environmental variation. The second kind of raptor searching and documentation is the *monitoring protocol*, which will be recommended for MSCP LRMP. This was based on which *survey* techniques were most useful, what has become standardized for raptors, and what will meet the objectives of a monitoring program (discussed below).

Based on a review of the MSCP Biological Monitoring Plan, discussions with the Contract Manager, and our knowledge of survey techniques that are widely accepted, we established guidelines for WRI biologists to follow for the breeding and wintering surveys (WRI 2004, Appendices A and B). As discussed in the Year 1 and 2 reports (WRI 2002, 2004), because of latitude, and the resulting mild climate of the MSCP area, raptor nesting activities can start as early as December and run into August. However, wintering raptors are commonly observed in this region December through February, with some remaining (or migrating through) into mid-March. Therefore, we have, somewhat arbitrarily, called field observations made December through February “winter” survey data. However, “breeding” season data are not limited to a specific timeframe, often overlap with the “winter” observation, and are based on observed behavior (e.g., copulation, nest building, incubation, bringing food to the nest, presence of young).

Table 1 provides a reminder of all the sites that were in the original list of those to be examined. One of the objectives of the 2003 fieldwork was to fill in some data gaps. We had difficulty gaining access to one site (San Diego National Wildlife Refuge/San Miguel Mountain, Site 23) because it involved the use of an access across private property. Table 1 does not reflect surveys that were conducted for the GE or numerous surveys conducted by WRI volunteers and cooperators. During this last year of study, we also continued our coordination with individuals responsible for managing the study sites to keep them apprised of project progress, maintain a point of contact, enlist their input, coordinate access, etc.

Although most of the fieldwork was conducted by vehicle and on foot, as described in WRI (2004, Appendices A and B), some observations, which were focused on the GE, were conducted by helicopter (WRI 2005).

RESULTS

LITERATURE REVIEW

The Project Bibliography has been completed (Appendix A); although, we would welcome any additions from those who review it. This bibliography is not intended to be comprehensive but is intended to provide the reader and local resource manager with important references that relate to: (1) relevant natural history of the target raptors; (2) the presence or distribution of the target raptors within the MSCP, and/or (3) survey or monitoring techniques that could be applied to the target raptor resources by land and wildlife managers within the MSCP. It is arranged by sections for each raptor target species, followed by a section on general raptor literature, with a focus on raptor management.

FIELD SURVEYS

The GE and the PF are addressed separately below because they are unique in both their biological status and their potential for being disturbed. The PF was only recently removed from the listing category and the GE has shown a marked (approximately 50 percent), and well-documented, decline in San Diego County.

Golden Eagle

The GE has been reported on separately (WRI 2005) for a number of reasons relating to resource protection. The detailed site-specific maps are provided in that document so that CDFG has the option of distributing those data separate from the other, less sensitive, raptor data depending on the recipient's need to know.

As an overview, however, after 16 years of consistent monitoring, we estimate that thirty one (31) pairs formerly occupied the San Diego MSCP. Today, fifteen (15) pairs are still active and sixteen (16) pairs have been extirpated. Most of these extirpations occurred in the last 35 years. The fifteen (15) breeding pairs of Golden Eagles remaining in the SD MSCP represent 30 percent of all the breeding Golden Eagles in San Diego County. Seven (7) of the fifteen (15) remaining active pairs within the SD MSCP are in serious jeopardy of being extirpated in the next 5-10 years. Three (3) of the seven (7) pairs predicted to become extirpated may, in fact, already be lost.

The first changes of significance that affected the SD MSCP Golden Eagle population were from intensive agriculture such as avocado and citrus groves. This agriculture replaced cattle grazing and grasslands. Some extirpations were documented to occur in San Diego County in the 1950s and 1960s, after the build-up of military personnel post-WWII, but most disappeared after the 1970s, when major freeways opened land for development that was formerly cattle ranches. Interstate and local freeways made access easy and allowed development to proceed.

Extirpated Golden Eagle territories were primarily located on private land (56 percent). Currently only three (20 percent) of the remaining pairs of Golden Eagles core nesting areas remain on private lands. Twelve (80 percent) of the currently active Golden Eagles within the SD MSCP nest on public land. *This is a significant and valuable opportunity for the future management and survival of Golden Eagles within the SD MSCP.*

In order to properly manage this far-ranging species, specific information about their ecological needs is required, including the limits of the core area around the nest, the primary foraging areas, and the limits of the defendable territory. These are provided in the Golden Eagle report (WRI 2005).

Peregrine Falcon

Breeding Season Results

Of the 12 current and/or historic PF territories known for the county, nine were (and, in five cases, are) located within the MSCP boundaries. Of the five territories located within the MSCP, only one territory is located at one of the study sites (Point Loma, Site 18; see Table 2). The status of that territory and others that we are aware of, within the MSCP, is as follows: Point Loma—active (likely produced young, 2002; was active, 2003); downtown San Diego—active (nest success not known, 2001-2003); La Jolla Cove—active (thought to have produced young, 2002); La Jolla Cliffs—active (nest success not known, 2001-2003); Downtown El Cajon—active (2002) but nest success not known.

Winter Results

A total of 14 PFs were documented during the winter months of 2002 and we believe this was typical for the study period (2001-2003). These were observed at ten study sites (Table 3). One individual was observed at each of nine sites, 2 at one site, and 3 were noted at, or near, another site (Point Loma; site 18). Most birds were observed along the coast or associated with large bodies of water, where shorebirds and other water-associated birds were abundant. Based on other observations, and input from knowledgeable raptor biologists, it is likely that there were roughly 20 PFs wintering in San Diego County during each of the period 2001-2003.

Other Raptors

Breeding Raptors

The raptor breeding season data, by study site, presented in Table 2 and Appendix B provides a picture of what each of the study sites can be expected to support under conditions of average-to-poor precipitation. Maps of all 45 study sites are provided. In cases where no data were collected, or data were combined between two sites, a note on the map provides that explanation. During the period 2001-2003, we examined 44 out of 45 sites (land access was not possible at SDNWR/San Miguel Mountain, Site 23 although we were able to survey a nearby GE nest by helicopter). We documented a total of 15 raptor species and 539 raptor breeding territories (excluding the CR but including 78 stick nests, which we could not positively identify as to raptor species). Of the 539 raptor breeding territories, 96 were target species (all but the BE, SH, and FH, which do not, currently, breed in the MSCP area). Sites varied greatly in their ability to support breeding raptors. Some sites didn't support more than one or two territories, while, others, like the Ramona Grasslands, supported almost 90 territories. Four sites supported no breeding raptors (see those with note "NBR"), while one site (Ramona Grasslands) supported 9 raptor species, including three target species.

The RT was the most commonly documented nesting raptor species, with a total of 177 nests and/or territories located on 34 sites. The next most commonly documented raptor

TABLE 2. Number of Raptor Nests and/or Territories by Site (2001-2003)

SPECIES**																					Stick Nest	Target Spp.	Total Spp.	Notes	Site No.	
SITE		AK	BE	BR	BO	CH	CR	FH	GE	GO	LO	NH	OS	PF	RS	RT	SO	SH	TV	WK						
1	Crestridge	1				1	2								3	2							1	9		1
2	Boden Canyon					2									2	2							2	6		2
3	Jamul Ranch					2	2			1					1	13					2	15	2	36		3
4	SDNWR*/Salt Works/Egger Ghio																						0	0	NBR	4
5	McGinty Mountain Complex					1									1	5							1	7		5
6	San Diego Bay NWR			1			1									1					2	0	5	Note 1	6	
7	Lake Hodges	1		2		2	1		1	2					5	8					1	3	23		7	
8	Penasquitos Lagoon																					0	0	Note 5	8	
9	Torrey Pines						6									1						0	7	Note 5	9	
10	Sycamore Canyon			1							3				1	1	1					0	7		10	
11	Iron Mountain			2		4	1		1	1			1		11	13			1	2		5	37	Note 2	11	
12	Otay Mountain						2		2						1	5					1	2	11		12	
13	Marron Valley						2		1							6					10	1	19		13	
14	Otay Lakes	1				2	2			1		2				4					1	4	4	17		14
15	SDNWR* Sweetwater Marsh											2				1						2	3	NBR	15	
16	San Vicente								1				3			2			2			1	8		16	
17	Sycuan Peak																					0	0	NBR	17	
18	Point Loma													1		1						1	2		18	
19	North Island				6								1			1					1	6	9		19	
20	Miramar Reservoir					1							1			1						1	3		20	
21	Mission Bay																					0	0	NBR	21	
22	Brown Field Complex	1			4		1					1				5					1	5	13		22	
23	SDNWR*/San Miguel Mountain								1													1	1	NSC	23	
24	Mission Trails					1									1	2					1	1	6		24	
25	Proctor Valley	3				1										3					1	1	8		25	
26	San Diego River	1				3	1		1	1					3	9						4	19		26	
27	Route 67 South/Iron Mtn #11																					0	0	Note 2	27	
28	San Dieguito Lagoon											1				4					1	1	6		28	
29	Route S-6																					0	0	Note 4	29	
30	Grasslands/Route 67	10			1	1	1		1	6					25	41			1	3		3	90		30	
31	Sloan Canyon						7		1						2	4					1	2	17		31	
32	Rockwood Canyon					1	1		1						1	4						2	8		32	
33	Penasquitos Canyon	3		2		7	1			2		2			9	4			1	6		9	37		33	
34	Hollenbeck Canyon	4				1	4					2			2	4					1	4	22		34	
35	Rock Mountain						1		1						1	2						1	5		35	
36	San Pasqual	1		4		2	2			3					9	16			1	2	7	2	47		36	
37	SDNWR*Tijuana Slough	1										2										2	3		37	
38	Route 94 (North and South)																					0	0	Note 3	38	
39	Immenschuh					1															1	1	2		39	
40	Los Montanas (North)														1	1					1	0	3		40	

TABLE 2. Number of Raptor Nests and/or Territories by Site (2001-2003)

SPECIES**																					Stick Nest	Target Spp.	Total Spp.	Notes	Site No.	
SITE		AK	BE	BR	BO	CH	CR	FH	GE	GO	LO	NH	OS	PF	RS	RT	SO	SH	TV	WK						
41	Los Montanas (South)															2							0	2		41
42	Rancho San Diego (East)	1				1										3					1	5	1	11		42
43	Rancho San Diego (West)					2									1							8	2	11		43
44	Border Fields	1		2		6				1		13			2	1					2	12	19	40		44
45	Sweetwater Reservoir					5	3			2					1	5					1	2	5	19		45
Total		29	0	14	11	47	41	0	12	20	3	25	6	1	83	177	1	0	6	25	78	96	579			

* San Diego National Wildlife Refuge.

NBR No breeding raptors observed.

NSC No formal raptor surveys conducted (see notes on topo report maps).

- (1) Breeding raptors and ravens observed in residential areas to east of study area.
- (2) Data for Route 67 South (# 27) and Iron Mountain (#11) were combined. See Iron Mountain (#11)).
- (3) The Route 94 transect overlaps other study sites. Data from this transect were assigned to other appropriate sites.
- (4) No data collected due to safety and access issues.
- (5) Data for Penasquitos Canyon (#8) combined with Torrey Pines (#9). See Torrey Pines (#9).

**Species:

AC American Crow	CR Common Raven	NH Northern Harrier	SO Screech Owl
AK American Kestrel	FH Ferruginous Hawk	OS Osprey	SS Sharp-shinned Hawk
BE Bald Eagle	GE Golden Eagle	PF Peregrine Falcon	SH Swainson's Hawk
BH Black Hawk	GO Great-horned Owl	PR Prairie Falcon	TV Turkey Vulture
BR Barn Owl	HH Harris' Hawk	RS Red-shouldered Hawk	WK White-tailed Kite
BO Burrowing Owl	LO Long-eared Owl	RT Red-tailed Hawk	
CH Cooper's Hawk	MR Merlin		

Table 3. Number* of raptors observed during the winter (primarily January, February, and December) surveys--2001-2003.

		SPECIES***																							Total	Total	Notes	Site
SITE		AK	BE	BR	BO	CH	CR	FH	GE	GO	LO	MR	NH	OS	PF	PR	RS	RT	SO	SS	SH	TV	WK	Target Spp.	Raptors		No.	
1	Crestridge					1	19										3	4				1		1	28		1	
2	Boden Canyon					2		2									2	6				3		4	15		2	
3	Jamul Ranch	5					6						1					7						1	19		3	
4	SDNWR*/Salt Works/Egger Ghio	2				1	2	1				1	4	3	1			4					1	7	20		4	
5	McGinty Mountain Complex						2										2	5						0	9		5	
6	San Diego Bay NWR	3					1	1					1	1	1									3	8		6	
7	Lake Hodges	5		4		3	2		2	2			1	2	1		8	36				2	3	7	71		7	
8	Penasquitos Lagoon																							0	0	Note 1	8	
9	Torrey Pines						12								2			2						2	16	Note 1	9	
10	Sycamore Canyon			2							6						2	2	2					0	14		10	
11	Iron Mountain			4		8	2		1	2				2			22	18				2	4	9	65	Note 2	11	
12	Otay Mountain	2					18		1				1					5						2	27		12	
13	Marron Valley	1				1	14										1	6				1		1	24		13	
14	Otay Lakes	5				3	10						2	2	1		1	5				1		6	30		14	
15	SDNWR* Sweetwater Marsh	2				1	1					1	3	1				5					1	4	15		15	
16	San Vicente																	4					2	0	6		16	
17	Sycuan Peak																							0	0	Note 3	17	
18	Point Loma	1				1	3								3			3						4	11	Note 4	18	
19	North Island	2			3		6							2				3						3	16	Notes 3 & 5	19	
20	Miramar Reservoir					2								2				2				2		2	8		20	
21	Mission Bay	2					2											2						0	6		21	
22	Brown Field Complex	4			3	1	8						4					7					2	8	29		22	
23	SDNWR*/San Miguel Mountain																							0	0	NWC	23	
24	Mission Trails	1				2	6										3	3				1	2	2	18		24	
25	Proctor Valley	3				1	132								1			8						2	145		25	
26	San Diego River	5				6	2			2			1	1			7	22					2	7	48		26	
27	Route 67 South/Iron Mtn #11																							0	0	Note 2	27	
28	San Dieguito Lagoon												2					8					2	2	12		28	
29	Route S-6	2																2						0	4	Note 4	29	
30	Grasslands/Route 67	7			2	1	3	9	3	6		1		1	1		4	12		1		40		16	91		30	
31	Sloan Canyon						5		1								2	1						1	9	Note 3	31	
32	Rockwood Canyon					2	2		2									6						4	12		32	
33	Penasquitos Canyon	6		4		14	2		4				4	2			18	8				2	12	22	76		33	
34	Hollenbeck Canyon	7				1	13		2				3			1	3	5						6	35		34	
35	Rock Mountain								3															3	3		35	
36	San Pasqual	11		7		2	6		2	6			1		1		16	57		1		3	8	6	121		36	
37	SDNWR*Tijuana Slough	3				1	4		1			1	2	1	1			4						5	18		37	
38	Route 94 (North and South)																							0	0	Note 6	38	
39	Immenschuh					1																		1	1		39	
40	Los Montanas (North)																	3						0	3		40	
41	Los Montanas (South)						4											3						0	7		41	
42	Rancho San Diego (East)	2				3	6											4					1	3	16		42	
43	Rancho San Diego (West)						3											1						0	4		43	
44	Border Fields	8											6				1	13					3	6	31		44	

Table 3. Number* of raptors observed during the winter (primarily January, February, and December) surveys--2001-2003.

45	Sweetwater Reservoir	9	2			3	38			2		1		1	1			5				1	1	6	64		45
Total		98	2	21	8	61	334	11	24	20	6	5	36	21	14	1	95	291	2	2	0	59	44	156	1155		

* Numbers refer to maximum number of birds observed.

** San Diego National Wildlife Refuge.

- (1) Data for Penasquitos Lagoon (#8) and Torrey Pines (#9) combined. See Torrey Pines (#9).
- (2) Data for Route 67 South (# 27) and Iron Mountain (#11) were combined. See Iron Mountain (#11).
- (3) Includes March survey.
- (4) Two PFs were observed to the north of this site, near the S.D. Airport.
- (5) Seven widely-spaced active burrows suggested that there were at least seven B0s on this study site.
- (6) The Route 94 transect overlaps other study sites. Data from this transect were assigned to other appropriate sites.

***Species

AK American Kestrel	CR Common Raven	MR Merlin	RS Red-shouldered Hawk	TV Turkey Vulture
BE Bald Eagle	FH Ferruginous Hawk	NH Northern Harrier	RT Red-tailed Hawk	WK White-tailed Kite
BR Barn Owl	GE Golden Eagle	OS Osprey	SO Screech Owl	
BO Burrowing Owl	GO Great-horned Owl	PF Peregrine Falcon	SS Sharp-shinned Hawk	
CH Cooper's Hawk	LO Long-eared Owl	PR Prairie Falcon	SH Swainson's Hawk	

nests/territories were those of the RS with 83 and the CH with 47. The CR (a non-raptor, but a species that can have an impact on raptors) was fourth in frequency with 41 nests/territories. The next level of frequency was shared by AK (29), NH (25), WK (25), and GO (20). To a great extent, this frequency distribution is a function of site size, amount of appropriate habitat, and sometimes local conditions on the respective sites.

Of the eight project target species, nesting was documented for five—CH, NH, GE, BO, and, PF. CH nesting was observed at the highest number of study sites, with nests and/or territories documented at 21 sites (48 percent of the 44 sites surveyed). GE was observed nesting at 11 sites (25 percent); while NH was documented at only 8 sites (18 percent) with 13 of the 25 territories found at Border Fields. BO were found nesting at only 3 (7 percent) of the sites and PF at only 1 (0.23 percent) of the sites.

The CH nested, primarily, at those sites that contain healthy riparian habitat; however, this species has become somewhat of a generalist and also nests elsewhere (see Discussion). GEs limited their nesting to sites with sheer cliffs away from human activity and close to nearby grasslands for hunting (see below). The NH and the PF were concentrated primarily along the coast. However, one PF pair attempted nesting in downtown El Cajon and a few scattered NHs were observed nesting at more inland sites. NHs nested in mostly coastal marsh and open field habitat; although we have observed NHs nesting in ruderal areas (J. Oakley, pers. comm.). PFs utilized mostly man-made structures, along the coast, with nearby sources of shorebirds and other prey. Most of BOs, located on the study sites, were found in sandy soil with low grass and open areas (see also WRI 2003, Lincer and Bloom 2003, in prep.). BE and FH winter within the MSCP but are not known to breed there. SHs only pass through during migration, are infrequently documented, and when they are, they are usually not within the MSCP. Some of the SH migrants seen are in the Ramona area and large numbers (over 5,200) have been recently documented migrating along the desert front to the east of the MSCP during the spring (Unitt 2004).

Based on the number of *all* nesting raptor species (plus the CR) and all the sites surveyed during the 2001-2003 breeding seasons, Site 30 (Ramona Grasslands/Route 67) contained the most nests/territories of all sites surveyed. Eighty-nine nests/territories were documented, representing nine raptor species (and 1 CR). The site to show the next highest number of territories was San Pasqual (Site 36) with 47 territories (including two CR and 7 unidentified stick nests that were not duplications of known territories). Border Fields State Park (Site 44) showed the next highest number of territories with 40 territories (including 12 non-duplicative unidentified stick nests).

Site 44 (Border Fields) contained the highest number of *target* species nests/territories of all sites surveyed (19). Penasquitos Canyon (Site 33) supported 9 target species territories while North Island (Site 19) supported 6 and Brown Field Complex (Site 22) and Iron Mountain (Site 11) tied, with both supporting 5 nests of the target raptor species.

Wintering Raptors

A total of 20 raptor species were documented on our study sites during the winter months (January, February, and December) of 2001-2003 (Table 3). Of course, at San Diego's latitude, a number of the resident breeders are actively nesting while many of the wintering birds are still on site. All target raptors, but the SH, were documented during the winter observation period (December-February). Numbers ranged from 0 to 22 individual target raptors per site for a total of 154 individuals for all study sites. Comparable numbers for all raptors (plus the Common Raven) were 0 to 145 as a range. A total of 1,153 wintering individuals were documented (or 819, without the ravens).

The CR was, clearly, the most common wintering bird of those surveyed for. The three most commonly documented wintering raptors were the RT, AK, and RS, with totals of 291, 98, and 95, respectively. Of those sites surveyed in this study, the following held the highest number of wintering individuals (raptors and ravens): Site 25 (Proctor Valley) – 145, Site 36 (San Pasqual) – 121, Site 30 (Ramona Grasslands) – 91 (which included 9-16 FHs; with 20 documented in 2005), Site 33 (Penasquitos Canyon) – 76, and Site 7 (Lake Hodges) – 71.

DISCUSSION

Weather as a Factor

In reviewing any body of data, it is important to consider how typical the sampling period was. So just how “typical” were 2001 through 2003? Drought plays a significant factor in the density and reproductive success of raptors and other predators. During the El Nino of 1998/99, NHs were breeding in areas where they have not bred since and in lower numbers in other locations. The demonstrable impacts of drought on GEs and Prairie Falcons, throughout southern California, were presented by Bittner et al. (2003). This study was conducted during the worst drought for San Diego in 160 years. This should be noted for future researchers.

Management and Enforcement Issues

Table 4 is a summary of management and enforcement issues by site. Clearly, some study sites are substantially impacted, either directly or indirectly, by human activities. Some sites are currently without major impacts. Unfortunately, many of the more diverse and potentially productive sites are the same ones that are experiencing multiple management and enforcement challenges. Of those that are obviously impacted, the following activities are the most common: humans walking or hiking (36 out of 45 sites or 80%) and pets, primarily dogs being allowed to run free, (26 out of 45 sites or 57 %).

Table 4. Management Enforcement Issues Identified by Raptor Study Site

Site No.	Name	Humans Walking/Hiking	Rock Climbing	Off-road Vehicle Use	Pets	Disking, etc. Agricultural Activities	Rodent/Ground Squirrel Poisoning	Construction/Development	Newly-developed Access Road(s)	Other
1	Crestridge	X								
2	Boden Canyon	X		X	X				X	6
3	Jamul Ranch									6?
4	SDNWR*/Salt Works	X								
5	McGinty Mountain Complex			X	X					
6	San Diego Bay NWR	X			X					
7	Lake Hodges	X	X	X	X			X	X	6
8	Penasquitos Lagoon	X			X					
9	Torrey Pines	X			X					7
10	Sycamore Canyon	X		X	X				X	
11	Iron Mountain	X	X	X	X				X	
12	Otay Mountain	?	X	X						1
13	Marron Valley	X	X	X	X					1
14	Otay Lakes	X			?				X	8
15	SDNWR* Sweetwater Marsh	X		X	X					
16	San Vicente	X	X		X				X	
17	Sycuan Peak									
18	Point Loma	X								
19	North Island	X								2
20	Miramar Reservoir	X						X		
21	Mission Bay	X		X	X			X		
22	Brown Field Complex	X		X	X			X		1,3,4
23	SDNWR*/San Miguel Mountain	X		X	X				X	
24	Mission Trails	X	X		X					
25	Proctor Valley	X		X	X		X	X		
26	San Diego River	X	X		X		X			7
27	Route 67 South	X		X				X	X	
28	San Dieguito Lagoon	X				X		X		
29	Route S-6	X						X		
30	Grasslands/Route 67	X	X		X	X	X	X	X	
31	Sloan Canyon	X			X					
32	Rockwood Canyon	X	X					X		
33	Penasquitos Canyon	X		X	X				X	
34	Hollenbeck Canyon									6
35	Rock Mountain	X	X							5
36	San Pasqual	X		X	X	X	X	X		5
37	SDNWR*Tijuana Slough	X		X	X	X	?			

Table 4. Management Enforcement Issues Identified by Raptor Study Site

38	Route 94 (North and South)									
39	Immenschuh									
40	Los Montanas (North)									
41	Los Montanas (South)									
42	Rancho San Diego (East)	X			X					
43	Rancho San Diego (West)	X			X					
44	Border Fields	X			X					1
45	Sweetwater Reservoir									

*San Diego National Wildlife Refuge

- (1) Border Patrol and illegal alien activities.
- (2) Conflicts with Navy goals and endangered species recovery program.
- (3) Potential conflict with future Navy goals at Satellite Surveillance Station.
- (4) Heavy predation by Coyotes and Barn owls.
- (5) Future threats from proposed trail construction and associated access to rock climbers, ORVs, etc. activities.
- (6) Shooting (legal and illegal).
- (7) Paragliding.
- (8) Cattle grazing.

Management Conflicts

The following are observed management conflicts, which lead to our recommended management and research (see Recommendations):

- As indicated above, human uses [rock-climbing, hiking, jogging, walking dogs (often without leashes), vehicular use, etc.] impact the normal behavior of raptors (and other wildlife).
- In many cases, the size of protected parcels is substantially smaller than that required by a raptor's functional territory, including foraging areas.
- The public/political pressure to create new trails into MSCP preserve lands provides a path for, and encourages, increased disturbance to raptors (and other wildlife).
- The public/political perception that MSCP preserve lands have been created primarily for active, and in some cases, consumptive, recreation, sets up an obvious conflict for managing raptors (and other wildlife).
- The constraint of using fire as a management tool in proximity to human habitation limits habitat management tools.
- Inadequate funding to both acquire important lands and properly manage MSCP lands which are acquired.

Raptor Monitoring

The following is a reiteration of considerations, regarding the MSCP Long-term Raptor Monitoring Program, that were presented previously (WRI 2004) and discussed elsewhere (Lincer and Bittner 2002; Lincer et al. 2003). For further reading, relevant issues are proposed and discussed by Oakley, Thomas, and Fancy (2003).

Sample Design

The ideal sample design should be:

1. Representative of the study area and the issues at hand. (e.g., habitat loss, disturbance, etc.) ;
2. Representative of the habitats of interest and the seasons during which those habitat support the monitored species (e.g., the MSCP not only provides important breeding habitat for numerous raptor but it is also a significant habitat for several wintering raptors, including some that are considered target raptors, like the PF, BE, FH, and BO);
3. Inclusive of all focus species or represent them in some functional way;
4. Sensitive to the objectives of the MSCP monitoring requirements;
5. Sensitive to logistics;
6. Statistically appropriate (which may be compromised by above logistics);
7. Able to predict, and take into consideration, *detectability* (i.e., how counts relate to the actual number of raptors in the sampled area; one approach is to use a "double count" approach). This objective may also be compromised by above logistics.

Questions to be Answered and Objectives to be Met

How will the data be used by the various management entities? When do they need what? An example of a clear monitoring objective would be, "Be able to detect a 25% change in population (individual species or overall raptor group?), in each chosen habitat, in 10 years." This is the approach that is being attempted by NARMS (North American Monitoring Strategy) but some of the best raptor monitoring minds are having a serious challenge addressing these objectives. It is entirely possible that we won't have enough observations for some species to detect a significant change in a timely manner.

Possible Monitoring Approaches

Levels of effort and *agency commitment* are, integrally tied. For instance, the MSCP program could adopt a:

1. Highly rigorous, scientific approach that would be costly but could withstand the most challenging statistical/legal tests, or
2. More practical, less expensive approach that would be more likely to be funded, and therefore carried out, but would stand the chance of being successfully, challenged at some time in the future.

As to *which, and how many, species* should be involved, the program could use a:

1. Multiple species approach, using selective target species only,
2. Multiple species approach, using selective target species, but recording all raptors (and ravens) observed,
3. Single species approach, using a keystone species, like the Golden Eagle or
4. Combination of the above.

Target Species and Other Multiple Species Approaches

A monitoring approach that focuses on one or more so-called "target" species has the appeal of apparent simplicity and the implication that these target species will, somehow, reflect a broader suite of species and be sensitive to whatever perturbations are experienced. Having surveyed raptors for many years, it is apparent that each species often responds to similar impacts differently. Although GOs and RTs might show similar population changes in response to small mammal population changes, and most raptors will show some response to a record-breaking drought, such as we have just experienced, there are likely more differences than similarities between species. Those differences are not only in *degree* but also in *direction*. For instance, GEs and PRs responded to the recent drought to different degrees (Bittner et al. 2003), with the PR being less impacted by presumed small mammal population decreases because it takes a wider range of prey species than the GE, which is heavily dependent on jackrabbit and ground squirrel populations. In addition, some raptors (e.g., GE) are far more negatively responsive to human activity than others (e.g., AKs, RTs, RSs, and some CHs). There are also differences in response, both within and between species, depending on the time of year (e.g., during the

breeding season vs. the wintering season) and where a disturbance occurs (e.g., on the hunting grounds or within the nest territory).

Regarding raptors responding in a different direction, one only needs to recognize that many different raptors require different habitats and, although not many species will persist if usable habitat is replaced with a development (although some CHs and RSs may defy this simplification), a conversion from one habitat/land use to another will often affect different species in different ways. For instance, if an extensive riparian habitat were to be replaced by an agricultural land use, and some hedge rows were to be left/created, we could expect that there would be a decrease in RSs, CHs, and several owl species. But, at the same time, there would likely be an increase in AKs, RTs, and perhaps WKs.

The point to the above exercise is that, if an arbitrary few species are chosen as “target” species, and the other raptors are not monitored, there will be a good chance that only some kinds of impacts will be reflected in the population trends of those raptors monitored. In our opinion, the MSCP Long-term Monitoring Program should include a broad-based approach, which documents all raptors observed and uses observed changes/trends to identify appropriate adaptive management strategies.

Single Species Monitoring Approach

Having sung the praises of a multiple raptor species approach (above), there is at least one raptor species in the western United States that has the ability to reflect regional trends in environmental health. This is the Golden Eagle. The attraction of using the GE, as a regional “miner’s canary,” is that (1) it requires a reasonably large and intact territory, and (2) there exists, in San Diego County, a unique and relevant historical regional database for this species. The Wildlife Research Institute has a long history of investigating the historical presence of GE in southern California, which includes the MSCP and environs (Bittner and Oakley 1999; WRI 2005). This collection of records has been compiled to reflect past documentation of GE pairs, their nesting success, hunting territories, and numbers of egg and /or young. The WRI database includes both active and extirpated territories beginning with records as early as 1864. WRI became involved in 1987 with the start of the San Diego GE Project (see Discussion in WRI 2005). *This project, in total, represents the longest such study of any eagle population in the Western Hemisphere, and is the second to longest in the world, next to one study in Switzerland.*

Providing this historical information, in conjunction with current trend data, is critical to managing the GE into the future. Only if we understand the extant population (within the context of the historical variation) can we properly evaluate the population and meet the needs of the species under current and future changing environmental and land-use conditions. If this is accomplished, it will reflect the success of the MSCP program.

RECOMMENDATIONS

Long-term MSCP Raptor Monitoring

Long-term monitoring is recommended under three categories: (1) Breeding Season, (2) Winter Season, and (3) Single Species Monitoring Program.

Breeding Season Monitoring Program

Twelve areas are recommended for breeding season portion of a Long-term Raptor Monitoring Program (Figure 2 and Table 5). Each Raptor Monitoring Area (RMA) consists of one to four of the individual raptor study sites that were surveyed during the period 2001-2003, the analysis of which led up to these recommendations. The choices of RMAs were based on a number of biological parameters (e.g., raptor diversity and population parameters, known history of raptor use), logistical considerations (how a monitor would move efficiently through a monitoring area), and a reasonable geographic coverage of the MSCP study area (see Methods). The Breeding Season Monitoring Program should, initially, be conducted every two years and encompass all 12 RMAs each time (i.e., don't conduct different portions of the total every other year). After a maximum of 5 monitoring events (i.e., 10 years), a statistical trend analysis should be conducted to determine if the frequency of every two years is adequate or, perhaps, unnecessarily frequent. Depending on the data, it may make sense to conduct this analysis earlier.

Raptor monitoring for the Breeding Season Monitoring Program should follow the protocol provided in Appendix C. This monitoring should be conducted by qualified raptor biologists with several years of relevant regional experience with the raptors found in the MSCP and proper training in the specific techniques necessary to conduct this monitoring.

Thanks to a grant from the San Diego Foundation, for post- (2003) fire studies, WRI was able to test this monitoring program on seven RMAs, representing varying degrees of being burned:

- B. Ramona Grasslands (Control Area)
- D. Iron Mountain (Burned)
- E. San Diego River (Burned)
- F. Sloan Canyon (Burned)
- H. Proctor Valley (Partially Burned)
- I. Rancho Jamul (Partially Burned)
- L. Otay Mountain (Burned)

The results of this monitoring effort were reported to the San Diego Natural History Museum (Lincer 2005).

Winter Season Monitoring Program

Because (1) the MSCP provides important wintering grounds for many raptors (some of which are *only* here during the winter), (2) coastal portions of the MSCP are not captured by the above breeding season monitoring approach, and (3) it is important to track at least three raptor species, that are primarily coastal in the MSCP, which have proven to be ideal bioindicators (PF, NH, and Osprey), we recommend conducting a winter monitoring program that focuses on the coastal portions of the MSCP (Figure 3). This, like the Breeding Season Monitoring program, should be conducted every two years (alternating years with the breeding season monitoring would be acceptable). After a maximum of 5 monitoring events (i.e., 10 years), a statistical trend analysis

should be conducted to determine if the frequency of every two years is adequate or, perhaps, unnecessarily frequent. Depending on the data, it may make sense to conduct this analysis earlier.

TABLE 5. Proposed MSCP Areas for Long-term Raptor Monitoring (Breeding Season)

<u>Area</u>	<u>Name</u>	<u>Study Sites* (original number(s))</u>
A	San Pasqual	San Pasqual (36), Lk. Hodges (7), Boden Cyn. (2), Rockwood (32)
B	Ramona Grasslands	Ramona Grasslands (30)
C	Penasquitos Canyon	Penasquitos Canyon (33)
D	Iron Mountain Complex	Iron Mountain**(11), San Vicente (16), Route 67 (27)
E	San Diego River	San Diego River (26)
F	Sloan Canyon	Sloan Canyon (31), McGinty Mtn. North (5), Sycuan Mtn. North (17)
G	Sweetwater River	Sweetwater Reservoir (45), Rcho. S.D. East (42), Rcho. S.D. West (43), San Miguel Mtn. North (23)
H	Proctor Valley	Proctor Valley (25), San Miguel Mtn. South (23), Upper Otay Lk.(14)
I	Rancho Jamul	Jamul Ranch (3), Hollenbeck Canyon (34)
J	Border Fields	Border Fields (44), Tijuana River (part)
K	Brown Field Complex	Brown Field (22), Otay River, Spring Cyn. (part), Dennery Cyn. (part)
L	Otay Mountain	Otay Mountain (12), Marron Valley (13), Lower Otay Lake (14)

* In some cases, only a portion of a study site is included because of access, visibility, or some other reason (see detailed maps, Appendix C, for details).

** Including Monte Vista Ranch.

Raptor monitoring for the Winter Season Monitoring Program should follow the protocol provided in Appendix C. This monitoring should be conducted from a vehicle, following the route depicted by Figure 3, and be conducted by qualified raptor biologists with several years of relevant regional experience with the raptors found in the MSCP.

Single Species Program

For the reasons covered in the Discussion section, we recommend that the GE (breeding season only) be used for the Single Species Program. Because of the dynamic nature of the GE pairs and the use of their territory, including their primary foraging area, these surveys should be conducted *every year* as they have been by WRI's biologists for the last 16 years. GE monitoring should follow the protocol that has been used for the San Diego GE Study for the last 16 years (Bittner and Oakley 1999, WRI 2005). WRI (2005) provides the details of both the breeding history of the GEs in the MSCP and recommendations on monitoring and future research. WRI (2005) is provided as a separate report for the protection and proper management of the GE. As an overview, observations must begin in December and go through June of each year. GEs begin courtship and nest building in December and January. They lay eggs in February and early March, hatch young in late March and April and fledge young in May and June. Therefore, it is essential that monitoring biologists be in the field for critical portions of the entire season (six months) to obtain all the data needed to monitor the GE population properly.

Aerial surveys have been a crucial part of the current study providing new insight into once-difficult areas to investigate potential territories. Patagial tags (and soon radio transmitters) placed on the GE's wings are now also an integral part of the eagle tracking process. Territory

Fig. 2. Prop'd RMAs (breeding)

Contact WRI for maps

integrity is fairly well documented in the San Diego MSCP and is being refined. See MSCP (2005) for more details.

Consistency in Monitoring

If data to be collected for this, or *any* monitoring program, are to have any utility in showing trends, they must be collected in a consistent fashion. As discussed above, the areas and routes to be monitored should be monitored frequently enough to reveal a complete picture of what is breeding and wintering on those respective areas and routes but these data are only a *sampling* of the entire MSCP. Therefore, it is extremely important that monitoring protocol is consistent both between sites/areas and over time (i.e., between years). To do this, a significant effort will have to go into selecting qualified raptor biologists, making sure that they are familiar with the required protocol, geography and species, and ensuring consistency between sites and years.

Other Recommendations

Management Needs and Enhancement Opportunities

- Restriction of inappropriate human activities where they are in conflict with, especially nesting, raptors.
- Apply the lessons learned in the development of the MSCP to the North and East County MSCPs and other HCPs.
- Develop a comprehensive management plan for the dwindling Burrowing Owl population within the MSCP.
- Selectively install artificial burrows, for BOs, and nest boxes for AKs, BRs, and Screech Owls (SOs). Keep in mind that BRs are an effective predator on not only small mammals but also medium size raptors, like the BO.
- Consider the use of grazing and/or fire as appropriate management tools to maintain grasslands, maintain/improve biological diversity, and manage fire fuel loading.

Recommended Research

- Transmitter study to better define the use of MSCP lands by GEs (initial studies in progress).
- Investigate the feasibility of reintroducing SHs into historical sites within the MSCP.
- Investigate the most efficient approaches to captive rearing and hacking BOs into appropriate habitat (either as is or as it can be modified and managed) within the MSCP.
- In order to prioritize the management of raptors that winter within the MSCP, but breed elsewhere (e.g., FH, MR, OS, BE, and some of the WK), determine the natal areas for these birds. If the natal areas have substantial threats, then no amount of MSCP management will have substantial positive impact.
- Document the growing OS population and determine emigration and immigration.
- Document the presence of, and habitat use by, crepuscular (BO) and nocturnal raptors (e.g., BR, SO, GO, Long-eared Owl).
- Document the recovery of raptors after the November 2003 fires and apply findings to future management strategies.

Fig. 3. Prop'd Winter Monit. Areas.

Contact WRI for Maps

ACKNOWLEDGEMENTS

We would like to thank and the CDFG NCCP Local Assistance Program for funding this project and the City of San Diego for funding Burrowing Owl work, upon which we drew to provide relevant data. David Mayer was a helpful and supportive Project Manager for CDFG. Holly Cheong (City of San Diego) was especially helpful in the set-up, management, and support of our Burrowing Owl efforts. Thanks go to WRI's Dave Bittner and Dr. Jim Hannan for reviewing an earlier version of this report and Dr. Richard Clark for his assistance in the literature search task. Phil Unitt (San Diego Natural History Museum) was very helpful by providing the database on raptor observations and associated maps from the ongoing County Bird Atlas Project. Many individuals, in numerous agencies and organizations provided assistance: Brian Collins, Monica Alfaro, Lorena Warner, Kate Shampaine, Clark Winchell (U.S. Fish and Wildlife Service); David Mayer, David Lawhead, Terri Stewart (State of California Department of Fish and Game); Robert Fisher, Mark Mendelsohn, Dr. Mark Fuller and Mike Kochert (USGS); Jeff Wells (U.S. Forest Service); Joe Caruso, Joe Weber (Otay Lakes City Water Supply); Dr. Joe Jehl (formerly with Hubbs-Sea World Research Institute); Charles Gailband (Chula Vista Nature Center); Pete Famolaro (Sweetwater Authority); Russell T. Donalson (Chugach Telecommunications & Computers, Inc.); Tamy Johniken, Dawn Larson, Steve Barnhill, Tammy Conkle (U.S. Navy); Brian R. Bonesteel, Billy Stewart (USDA-APHIS-Wildlife Services); Dr. Geoff Holroyd, Dr. Troy Wellicome, Ms. Helen Trefry (Canadian Wildlife Service); Richard Barber, Robert Jones (Palomar Audubon Society); Dieter Bothe (Pardee Homes); Jack Barclay (Albion Environmental); Scott Taylor (HELIX Environmental Planning); Christina Schaefer (EDAW); Pete Bloom (Western Foundation for Vertebrate Zoology); Dr. Tom Scott, Kristine Preston (U.C. Riverside); Dr. Jon Bart (Boise State University); Dr. Noel Snyder; Mary Ann McLeod.; Dr. Dan Varland; and Richard Glinski. We would like to thank the following individuals for sharing their historic data on the GE and their nest sites: John Oakley, Co-Director, WRI Eagle Project; J. B. Dixon, A.M. Ingersoll, Harry L. Heaton, Raymond Quigley, E.E. SeChrist, Maurice Burns, N. K. & B.P. Carpenter, Craig Culver, and Ed N. Harrison. Many thanks also go to the volunteers of the WRI Eagle Project and WRI's Hawk Watch. Finally, we recognize and appreciate that the following individuals have volunteered endless hour of field time: Dave Bittner, John Oakley, Jeff Wells, John Colton, Jeff Lincer, Tom Scott, Chris Meador, Jim Hannan, Nick Muscolino, Randy West, Dave Seals, Kate Shampaine, Geoff Rodgers, Jim Bryan, Craig Culver, Paul Jorgenson, and many others.

LITERATURE CITED

- Bittner, J.D. and J. Oakley. 1999. Status of Golden Eagles in Southern California. Raptor Research Foundation Conference. Golden Eagle Symposium. November 2. La Paz, Baja California.
- Bittner, J.D., J.L. Lincer, J. Oakley, Nick Muscolino, and J. Hannan. 2003. Golden Eagle reproduction in a drought period. Raptor Research Foundation's Annual Scientific Conference. September 2-7. Anchorage, AK.
- County of San Diego. 1997. Multiple Species Conservation Program; County of San Diego, Subarea Plan. Adopted by the Board of Supervisors on October 22, 1997.
- Hurlburt, T. 1971. [Probability of Interspecific Encounter]. In Gotelli, N.J. and G.L. Entsminger. 2001. EcoSim: no model software for ecology. Version 7.0. Acquired Intelligence, Inc. and Kesey-Bear. HTTP: [\\Homepage.together.net\gentsmin\ecosim\ecosim.htm](http://Homepage.together.net/gentsmin/ecosim/ecosim.htm).
- Lincer, J. 2005. Post-Fire Raptor Monitoring Report. Letter report to Dr. Mick Hager, Executive Director, San Diego Natural History Museum. 20 January. Prepared for the San Diego Foundation.
- Lincer, J.L. and P.H. Bloom. 2003. The status of the burrowing owl (*Athene cunicularia*) in San Diego County, CA. California Burrowing Owl Symposium. 11-12 November. Sacramento, CA.
- Lincer, J.L. and P.H. Bloom. In prep. The status of the burrowing owl (*Athene cunicularia*) in San Diego County, CA. Proceedings of the California Burrowing Owl Symposium. 11-12 November. Sacramento, CA.
- Lincer, J.L. and J.D. Bittner. 2002. Use of Raptors to monitor Habitat Conservation Plans. Poster presented at the Raptor Research Foundation's Annual Scientific Conference. September 24-28. New Orleans, Louisiana.
- Lincer, J.L., J.D. Bittner, N. Muscolino, and, L. Swartz. 2003. A Raptor Protocol for Monitoring HCPs. Paper to be presented at the Raptor Research Foundation's Annual Scientific Conference. September 2-7. Anchorage, AK.
- Ogden. 1996. Biological Monitoring for the Multiple Species Conservation Program. Prepared for the City of San Diego, California Department of Fish and Game, and U.S. Fish and Wildlife Service. Revised April 25.
- Oakley, K.L., L.P. Thomas, and S.G. Fancy. 2003. Guidelines for long-term monitoring protocols. Wildlife Society Bull., 31 (4): 1000-1003.
- WRI (Wildlife Research Institute, Inc.). 2002. Year 1 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2001). Prepared for California Department of Fish and Game. 30 July.

WRI (Wildlife Research Institute, Inc.). 2003. Burrowing Owl Surveys on City of San Diego Properties (for the period January 1, 2001 – March 24, 2003). Prepared for the City of San Diego, Planning Department. 24 March.

WRI (Wildlife Research Institute, Inc.). 2005. Final Report for NCCP/MSCP Raptor Monitoring Project-Golden Eagles of the San Diego Multiple Species Conservation Plan Area 2001-2003. Prepared for California Department of Fish and Game. 31 March.

APPENDIX A

PROJECT BIBLIOGRAPHY

Bald Eagle

- Anderson, D.W. and J.J. Hickey. 1972. Eggshell changes in certain North American birds. Proc. Int. Ornithol. Congr. 15:514-540.
- Buehler, D.A. 2000. Bald Eagle *Haliaeetus leucocephalus*. In The Birds of North America, No. 506. (A. Poole and F. Gill, eds.). The Academy of Natural Sciences, Philadelphia, PA and The American Ornithologists' Union, Washington, D.C. 40 pp.
- Buehler, D.A., T.J. Mersmann, J.D. Fraser, and J.K.D. Seegar. 1991. Effects of human activity on Bald Eagle distribution on the northern Chesapeake Bay. J. Wildl. Manage. 55(2):282-290.
- California. Department of Fish and Game. 1980. Bald Eagle *Haliaeetus leucocephalus* At the Crossroads. 89-90 pp.
- California. Department of Fish and Game. 1982. Bald Eagle. *Haliaeetus leucocephalus* California's Wildlife. 2:122 p.
- Dawson, W.L. 1923. The Southern Bald Eagle. In The Birds of California, Vol. 3. South Moulton Company, San Diego, Calif. Student's ed. 1712-1717 pp.
- Detrich, P.J., D.K. Garcelon. 1986. Criteria and habitat evaluation for Bald Eagle reintroduction in coastal California. The Resources Agency Department of Fish and Game. 1-32 pp.
- Detrich, P. J. 1981. Historic range of breeding Bald Eagles in California. Unpubl. Report., Redding, California. California State University, Chico 17 pp.
- Detrich, P.J. 1986. Status and distribution of the Bald Eagle in California. MS Thesis, California State University, Chico 112 pp.
- Forbis, L.A., B. Johnston, A. M. Camarena, and D. McKinney. 1977. Bald Eagle-habitat management guidelines. U.S.D.A. Forest Service, California Region. 60 pp.
- Gerrard, J.M. and G.R. Bortolotti. 1998. The Bald Eagle: haunts and habits of a wilderness monarch. Smithsonian Inst. Press, Washington, D.C.
- Green, N. 1985. The Bald Eagle. Audubon Wildlife Report. 508-529 pp.
- Grinnell, J., and A. H. Miller. 1944. The distribution of the birds of California. *Pacific Coast Avifauna* 27:17, 106.

- Harmata, A.R. 1993. Heavy metal and pesticide contamination of Bald and Golden Eagles in the western United States. Unpubl. USEPA, December 1993. 43 pp.
- Hastings, B. and C. Comp. 1988. Midwinter Bald Eagle survey report. National Wildlife Federation, Washington, D.C., 1986-1988.
- Henny, C.J. and R.G. Anthony. 1989. Bald Eagle And Osprey. Western Raptor Mgmt. Symposium and Workshop. 12:317.
- Howell, A.H. and F.L. Jaques. 1932. Florida bird life. Florida Department of Game and Fresh Water Fish. 182-183 pp.
- Institute for Wildlife Research. 1989. Midwinter Bald Eagle survey report. National Wildlife Federation, Washington, D.C.
- Jacobson, S.L. 1987. Bald Eagle habitat capability model. [Second draft.]. Unpubl. Report. U.S. Forest Service, Shasta-Trinity National Forest, Redding, California. 35 pp.
- Jones and Stokes Associates, 1982. Investigation of wintering Bald Eagles at Lake Mathews, Riverside County, California. Jones & Stokes, Inc., Sacramento, CA. (Prepared for the Metropolitan Water District of Southern California)
- Jurek, R.M. 1979. Southern Bald Eagle. Job Progress Report. State of California The Resource Agency, Department of Fish and Game.
- Jurek, R.M. 1982. Endangered, threatened and rare wildlife. Job Progress Report. State of California The Resource Agency, Department of Fish and Game.
- Jurek, R.M. 1982. Endangered, threatened and rare wildlife. Job Final Report. State of California, The Resource Agency, Department of Fish and Game.
- Jurek, R.M. 1988. Five year status report Bald Eagle. State of California The Resource Agency Department of Fish and Game. 1-15 pp.
- Jurek, R.M. 1990. California Bald Eagle breeding population survey and trend, 1970-90. State of California, Dept. of Fish and Game. 16 pp.
- Jurek, R.M., D.M. Hom, and C. Roberts 1986. California mid winter Bald Eagle survey. State of California, The Resource Agency, Department of Fish and Game. 4pp.
- Knight, R.L. 1984. Responses of wintering Bald Eagles to boating activity. *J. Wildl. Manage.* 48(3): 999-1004 pp..
- Lehman, R.N. 1979. A survey of selected habitat features of 95 Bald Eagle nest sites in California. Wildl. Management Branch Administrative Report 79-1:1-23
- Lehman, R.N. 1981. Breeding status and management of Bald Eagles in California. State of California The Resources Agency, Department. of Fish and Game. 83-1:1-24

- Lincer, J.L. 1981. "Bald Eagle Management at the Local Government Level." Paper presented at the 45th Annual Meeting of the Florida Academy of Sciences, April 30-May 2, Orlando, FL. *Florida Scientist*, 44 (1): 36-37.
- Lincer, J.L. 1982. "Bald Eagle: Symbol of Symbols." Invited editorial for ENFO Newsletter (a publication of the Florida Conservation Foundation), Vol. 82, No. 3. Winter Park, FL.
- Lincer, J.L. 1982. "Protecting Endangered Species at the Local Governmental Level." Paper presented at the 46th Annual Meeting of the Florida Academy of Sciences, April 22-24, DeLand, FL. *Florida Scientist*, 45(1): 40.
- Lincer, J.L. (Consulting Editor). 1989. Raptor Habitat Management Under the U.S. Bureau of Land Management Multiple-Use Mandate by R.R. Olendorff *et al.* Raptor Research Report No. 8, Raptor Research Foundation, Inc. Allen Press. 80 pp.
- Lincer, J.L., R.G. Brooks and B.L. Valla. 1991. "Managing Bald Eagles at the Local Level: A Prototypical Ordinance." Presented by J.L. Lincer at the Raptor Research Foundation Conference, November 6-9, Tulsa, OK.
- Lincer, J.L., W.S. Clark and M. Le Franc. 1979. Working Bibliography of the Bald Eagle. A comprehensive guide to the literature on the bald eagle. 2,000 refs. 268 pages with permuted keyword sort for index. National Wildlife Federation, Washington, D.C.
- Lincer, J.L., B. Millsap and G. Holder. 1988. "Bald Eagle Buffer Zones: Do They Work in Florida?" Presented at Raptor Research Foundation Annual Meeting, October 26-29, Minneapolis, MN.
- McWilliams, S.R., J.P. Dunn, and D.G. Raveling. 1994. Predator-prey interactions between eagles and cackling Canada and Ross' Geese during winter in California. *Wilson Bull.* 106:272-288.
- National Wildlife Federation. 1989. Proceedings of the Western Raptor Management Symposium and Workshop. National Wildlife Federation, Washington, D.C., Series No. 12. October 26-28, 1987, Boise, Idaho.
- Simmons, T., S. K. Sherrod, M.W. Collopy and M. A. Jenkins. 1988. Restoring the Bald Eagle. *American Scientist* 76(3): 252-260.
- Smith, B. 1989. Plan for Bald Eagles sought at Cachuma. Santa Barbara News Press (Santa Barbara, CA) (March 6):A1-A5.
- Solomon, S. and T. Newlon. 1991. Living with eagles. Status report and recommendations. Northwest Renewable Resources Center. 9-47 pp.
- Stalmaster, M.V. 1987. The Bald Eagle. Universe Books, New York.

- Stalmaster, M.V. and J.R. Newman. 1978. Behavioral responses of wintering Bald Eagles to human activity. *J. Wildl. Manage.* 42(3): 506-513.
- Steenhof, K. 1988. Identifying potential Bald Eagle nesting habitat; a review of the state of the art. pp. 31-59. *In* D. K. Garcelon and G. W. Roemer [eds.], Proceedings of the International Symposium on Raptor Reintroduction 1985. Institute for Wildlife Studies, Arcata, CA.
- Steinhart, P. 1990. Bald Eagle. *Haliaeetus leucocephalus* California. Wildlife. heritage: threatened and endangered animals in the golden state California Department of Fish and Game. 15-17 pp.
- Thompson, R.A. 1973. Bald Eagle nesting surveys in California. United States Fish and Wildlife Service. 48 pp.
- U.S Fish and Wildlife. 1994. Federal Register. Reclassify the Bald Eagle from endangered to threatened in most of the lower 48 states. 59: 132.
- Wood, P.B., D.A. Buehler, and M.A. Byrd. 1990. Raptor status report-Bald Eagle. *In* Proceedings of the southeast raptor management symposium and workshop (B. Giron Pendleton, ed.). National Wildlife Federation. Washington, D.C. Pp. 13-21.
- WRI (Wildlife Research Institute, Inc.). 2002. Year 1 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2001). Prepared for California Department of Fish and Game. 30 July.
- WRI (Wildlife Research Institute, Inc.). 2004. Year 2 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2002). Prepared for California Department of Fish and Game. 19 March.
- WRI (Wildlife Research Institute, Inc.). 2005. Final Report – NCCP/MSCP Raptor Monitoring Project (2001-2003). Prepared for California Department of Fish and Game. 31 March.

Burrowing Owl

- Allaback, M., J. Barclay. 1994. Burrowing Owl *Speotyto (Athene) cunicularia* Reference Manual. Proceedings of Burrowing Owl consortium of August 1993.
- Anderson, S.H., L.W. Ayers, J.A. Dechant, M.T. Green, W.H. Howe, S.L. Jones, D.S. Klute, D.K. Rosenberg, S.R. Sheffield, and T.S. Zimmerman. 2001. Status assessment and conservation plan for the western Burrowing Owl in the United States. Administrative Report. U.S. Department of Interior, Fish and Wildlife Service, Denver, CO.
- Banuelos, G. 1997. The one-way door trap: an alternative trapping technique for Burrowing Owls. *J. Raptor Res.* 9:122-124.

- Barclay, J., D. Plumpton and B. Walton 1998. (The California Burrowing Owl consortium) Burrowing Owl conservation in California: issues and challenges. Poster presented at Second International Burrowing Owl Symposium. The Raptor Research Foundation, Inc. Ogden, UT.
- Benson, P.C. 1981. Large raptor electrocution and power pole utilization: a study in six western states. Ph.D. thesis, Brigham Young University, Provo, UT.
- Bent, A.C. 1938. Life histories of North American birds of prey. U.S. Natl. Mus. Bull. No. 170.
- Best, T.R. 1969. Habitat, annual cycle, and food of burrowing owls in southwestern New Mexico. M.S. Thesis, New Mexico State Univ., Las Cruces. 34pp.
- Bloom, P.H. 1994. Ramona Airport Spring Raptor Census Report (*in* BFMA. 1997).
- BFMA (Brian F. Mooney Associates). 1997. Ramona Airport, Airport Master Plan Improvements, Ramona, CA. Vol. 1. Prepared for the County of San Diego and the U.S. Department of Transportation. March.
- Boggs, D.F. 1980. Respiratory adaptations of a burrow-dwelling bird to hypoxia and hypercarbia. Ph.D. thesis, University of Montana.
- Buchanan, J.T. 1997. A spatial analysis of the Burrowing Owl (*Speotyto cunicularia*) population in Santa Clara County, California, using a geographic information system. pp. 90-96. In Biology and conservation of owls of the northern hemisphere; second international symposium. February 5-9, 1997. Winnipeg, Manitoba, Canada, (J.R. Duncan, D.H. Johnson, and T.H. Nicholls, eds.). U.S. Department of Agriculture. Forest Service. North Central Forest Experiment Station, St. Paul, MN.
- Burrows, C.W. 1989. Diets of five species of desert owls. *Western Birds*. 20:1-10.
- Butts, K.O. 1973. Life History and Habitat Requirements of Burrowing Owls in Western Oklahoma Master's Thesis. Oklahoma State University, Stillwater. 188 Pp.
- CBOC (California Burrowing Owl Consortium). 1997. Burrowing Owl Survey Protocol and Mitigation Guidelines. *In* Lincer and Steenhof. 1997. The Burrowing Owl, its Biology and Management. *Raptor Res. Report* 9: Appendix B.
- CDFG (California Department of Fish and Game). 1995. Staff Report on Burrowing Owl Mitigation. Memorandum to Division Chiefs and Regional Managers from C.F. Raysbrook, Interim Director, California Department of Fish and Game. October 17.
- Clark, Greg. 2001. Burrowing Owl artificial nest box project an Arizona partners in flight habitat substitution project. Website citation: www.mirror-pole.com/burr_owl/bur_owl1.html.

- Clark, R.J., J.L. Lincer, and J.S. Clark. 1997. A Bibliography on the Burrowing Owl (*Speotyto cunicularia*). In *The Burrowing Owl, Its Biology and Management*. Proceedings of the First International Burrowing Owl Symposium. Raptor Research Reports No. 9:145-170.
- Climpson, J.T. 1977. Feeding ecology and selected other aspects of the behavior and ecology of the Burrowing Owl (*Speotyto cunicularia*). M.S. thesis, Washington State University, Pullman, WA.
- Colvée, Salvador. 1996. Ecología alimentaria del Mochuelo de Hoyos (*Athene cunicularia*) en la Peninsula de Paraguana. MS. Dissertation. Universidad Simon Bolivar. 34 pp.
- Coulombe, H.N. 1968. Energy exchange in the biology of the Western Burrowing Owl, *Speotyto cunicularia*. Ph.D. thesis. University of California at Los Angeles, CA.
- Coulombe, H.N. 1971. Behavior and population ecology of the Burrowing Owl, *Speotyto cunicularia*, in the Imperial Valley of California. *Condor*. 73:162-176.
- Delevoryas, P. 1997. Relocation of Burrowing Owls during courtship period. In Lincer and Steenhof (Eds.). 1997. *The Burrowing Owl, its Biology and Management Raptor Res. Reports*, 9:138-144.
- DeSante, D.F., E.D. Ruhlen, S. L. Adamany, K.M. Burton, and S. Amin. 1997. A census of burrowing owls in Central California in 1991. In Lincer, J. L. and K. Steenhof. 1997. *The Burrowing Owl, Its Biology and Management. Raptor Research Report Number* 9:38-48.
- DeSante, D.F., E.D. Ruhlen, and D.K. Rosenberg. 2002. Density and Abundance of Burrowing Owls in the agricultural matrix of the Imperial Valley, California. *Press, Studies in Avian Biology*. 9 pp.
- Drost, C. A and R. C. McCluskey. 1992. Extirpation of alternative prey during a small rodent crash. *Oecologia* (Berlin) 92(2):301-304
- Duxbury J.M. and G.L. Holroyd. 1995. A standardized, roadside Burrowing Owl survey technique. Canadian Wildlife Service, Environment Canada. Department of Renewable Resources. 7 pp.
- Duxbury, J.M. and G.L. Holroyd. 1998. A Standardized, Roadside Burrowing Owl Survey Technique. Paper presented at The Second International Burrowing Owl Symposium held in conjunction with the 1998 Annual Raptor Research Foundation Meeting. 29-30 September. Ogden, UT.
- Holroyd, G.L. and T.I. Wellicome. 1998. Report on the Burrowing Owl Conservation Workshop. Pages 612-615 in Duncan, J.R., D.H. Johnson and T.H. Nicholls (eds.). *Biology and Conservation of Owls of the Northern Hemisphere, Second International Symposium, February 5-9, 1997, Winnipeg, Manitoba, Canada*

- EDAW. 2001. Wildlife Biological Technical Report for the East Otay Mesa Specific Plan Amendment Area, San Diego, California. Prepared for County of San Diego Department of Planning and Land Use. October.
- Feeney, L.R. 1997. Burrowing Owl site tenacity associated with relocation efforts. *J. Raptor Res.* 9:132-137.
- Gervais, J.A., D.K. Rosenberg, and R.G. Anthony. 2001. Burrowing Owl space use and pesticide exposure risk in an agricultural landscape. *J. Wildl. Manage.* 31 pp.
- Green, G. A., R. E., Fitzner, R. G Anthony, L E Rogers. 1993. Comparative diets of burrowing owls in Oregon and Washington. *Northwest Science* 67(2): 88-93.
- Grinnell, J. and A.H. Miller. 1944. The distribution of the birds of California. Contribution from the Museum of Vertebrate Zoology of the University of California. Reprinted by Artemisia Press. Lee Vining, CA. 617 pp.
- Haley, K.L. 2002. The role of food limitation and predation on reproductive success of Burrowing Owls in Southern California. (Thesis) submitted to Oregon State University. 47 pp.
- Halverson, W.S. and A.C. Crabb. (Eds). 1994. Natural history and protection of Burrowing Owls. Proceedings of the Sixteenth Vertebrate Pest Conference. March 1-3, 1994. Santa Clara, California. Published at University of California, Davis. 83-86.
- Haug, E.A. and A.B. Didiuk. 1981. Use of recorded calls to detect Burrowing Owls. *J. Field Ornithol.* 64(2):188-194.
- Haug, E.A., B.A. Millsap, and M.S. Martell. 1993. Burrowing Owl (*Speotyto cunicularia*). In The Birds of North America, No. 61:2-20. (A. Poole and F. Gill, eds.). The Academy of Natural Sciences, Philadelphia, PA and The American Ornithologists' Union, Washington, D.C.
- HELIX Environmental Planning, Inc. 1999. San Diego Air Commerce Center at Brown Field Airport Master Plan Biological Assessment. United States Department of Transportation Federal Aviation Administration. 20 pp.
- HELIX Environmental Planning, Inc. 2001. Draft EIR/EIS of Route 905. Prepared for Federal Highway Administration and Caltrans.
- Hennings, L.T. 1970. Life history of the Burrowing Owl at the Oakland Airport, Alameda County, California. MS. Dissertation, University of California, Berkeley.
- Henny, C.J. and L.J. Blus. 1981. Artificial burrows provide new insight into Burrowing Owl nesting biology. *Raptor Res.* 15:82-85.

- Holroyd, G.L. and H. Trefry. 1998. Migration and winter biology of Burrowing Owls in the USA. *In* Second international Burrowing Owl symposium. The Raptor Research Foundation, Inc. 3 pp.
- Holroyd, G.L., R. Rodriguez-Estrella, and S.R. Sheffield. 2001. Conservation of the Burrowing Owl in Western North America: issues, challenges, and recommendations. *J. Raptor Res.* 35(4):399-407.
- Holroyd, G.L. 1998. A Burrowing Owl conservation action plan – what should it contain? *In* Second international Burrowing Owl symposium. The Raptor Research Foundation, Inc. 5 pp.
- Hunting, K. 1998. Mitigating impacts to Burrowing Owl populations: case studies in California. *In* Second international Burrowing Owl symposium. The Raptor Research Foundation, Inc. 4 pp.
- James, P.C. and R.H.M. Espie. 1997. Current status of the Burrowing Owl in North America: an agency survey. *J. Raptor Res.* 9:3-5.
- James, P.C., T.J. Ethier, and M.K. Toutloff. 1997. Parameters of a declining burrowing owl population in Saskatchewan. *In* Lincer and Steenhof (Eds.). 1997. The Burrowing Owl, its Biology and Management *Raptor Res. Reports*, 9:34-37.
- Johnson, B.S. 1988. Viability of small populations of the burrowing owl. *Bull. Ecol. Soc. Amer.* Suppl. 69 (2):182.
- Johnson, B.S. 1992. Effects of demography, mating patterns, and sociality on the population genetics and dynamics of the Burrowing Owl *Athene cunicularia*. Doctoral Dissertation. University of California, Davis.
- Johnson, B.S. 1997. Characterization of population family genetics of the burrowing owl by DNA fingerprinting with pV47-2. *In* Lincer and Steenhof (Eds.). 1997. The Burrowing Owl, its Biology and Management *Raptor Res. Reports*, 9:58-63.
- Landry, R.E. 1979. Growth and Development of the Burrowing Owl, *Athene Cunicularia*. Long Beach, California State University. MA Dissertation. 74 pp
- Lehman, R.N., L.B. Carpenter, K. Steenhof, and M.N. Kochert. 1991. Assessing relative abundance and reproductive success of shrub steppe raptors. *J. Field Ornithol.* 69(2):244-256.
- Lincer, J.L. 1997. Toward an action plan: *In* Lincer and Steenhof (Eds.). 1997. The Burrowing Owl, its Biology and Management *Raptor Res. Reports*, 9:5:11-13.
- Lincer, J.L. and K. Steenhof (Eds.). 1997. The Burrowing Owl, its biology and management including the proceedings of the first international Burrowing Owl symposium. *Raptor Res. Rep.* 9:1-177.

- Lincer, J.L., R.J. Clark and J.S. Clark. 1998. Toward an update on Clark, Lincer and Clark's bibliography on *Speotyto cunicularia*. Presented at the second international Burrowing Owl symposium, Ogden, Utah. September 29-30, 1998.
- Lincer, J.L., R.J. Clark and J.S. Clark. 1998 Towards a Burrowing Owl bibliography. *In* Second international Burrowing Owl symposium. The Raptor Research Foundation, Inc. 2 pp.
- Martell, M.S. 1991. Grassland owls. Pp. 96-104 in Proceedings of the Midwest raptor management symposium and workshop (B.G. Pendleton and D.L. Krahe, Eds.). National Wildlife Fed. Sci. Tech. Series, No. 15. Washington, D.C.
- MacCracken, J.G., D.W. Uresk, and R.M. Hansen. 1985. Vegetation and soils of burrowing owl nest sites in Conata Basin, South Dakota. *Condor* 87:152-154.
- Merkel, K.W. and D.A. Mayer. 2000. 1999 Maintenance and monitoring report for the artificially created burrows for the Burrowing Owl (*Speotyto cunicularia*) at the Otay Water District use area property, Chula Vista, California. Merkel & Associates, Inc. 99-046-01.
- Millsap, B.A., M.I. Bellocq, and M. Mullenix. 1997. Overview of literature on the burrowing owl. *In* Lincer and Steenhof (Eds.). 1997. The Burrowing Owl, its Biology and Management *Raptor Res. Reports*, 9:6-10.
- Ogden (Ogden Environmental and Energy Services Co.). 1992. Otay Ranch Raptor Management Study. Submitted to Otay Ranch Project Team, Chula Vista, California.
- Ogden. 1996. Biological Monitoring for the Multiple Species Conservation Program. Prepared for the City of San Diego, California Department of Fish and Game, and U.S. Fish and Wildlife Service. Revised 25 April.
- Palacios, E., D.W. Anderson, E. Mellink, and S. Gonzalez-Guzman. 2000. Distribution and abundance of Burrowing Owls on the peninsula and islands of Baja California. *Western Birds*. 31:89-99.
- Poulin, R.G., T. Wellicome, R. Longmuir, and Dave Scobie. No date. Burrowing Owl nest box: construction and installation procedures. Saskatchewan Environment and Resource Management. Fish and Wildlife Branch.
- Priest, J.E. 1997. Age identification of nestling Burrowing Owls. *J. Raptor Res.* 9:125-127.
- Recon. 2001. Otay Mesa Generating Project Biological Resources Mitigation Implementation and Monitoring Plan. Prepared for Otay Mesa Generating Company. 5 September. (Note: 2002-2003 updates on burrowing owl distribution based on personal communications between Fred Edwards and David Mayer, CDFG.)
- Raptor Research Foundation, Inc. 2001. The Second International Burrowing Owl Symposium. *J. Raptor Res.* 35(4): 269-417.

- Rich, T. 1984. Monitoring burrowing owl populations: implications of burrow re-use. *Wildl. Soc. Bull.* 12:178-180.
- Rosenberg, D.K. and K.L.Haley. 2000. The ecology of Burrowing Owls in the agroecosystem of the Imperial Valley, California.. *In Press, Studies in Avian Biology.* 40 pp.
- Rosier, J.R., N.A. Ronan, and D.K. Rosenberg. 2001. Breeding season survival and dispersal of Burrowing Owls in an extensive California grassland. *Review: Condor.* 16 pp.
- Rowe, Matthew P. 1984. California ground squirrels and their burrow coinhabitants: communicatory coevolution between predators and prey. Ph.d. Dissertation. Davis, University of California. 110 pp
- Sheffield, S.R. 1998. Conservation of the Burrowing Owl in north America: problems, issues, and solutions. *In Second international Burrowing Owl symposium.* The Raptor Research Foundation, Inc. 5 pp.
- Smith, B.W. and J.R. Belthoff. 1998. Burrowing Owls and development: results of short-distance nest burrow relocations to avoid construction impacts. *In Second international Burrowing Owl symposium.* The Raptor Research Foundation, Inc. 4 pp.
- Smith, B.W. and J.R. Belthoff. 1999. Ectoparasites on Burrowing Owls: Potential effects on nest-site reuse and growth, body condition, and survival of juveniles. *In Raptor Research Foundation Annual Meeting.* November 3-7, 1999, La Paz, Baja California Sur, Mexico 12 pp.
- Smith, B.W. and J.R. Belthoff. 2001. Burrowing Owls and development: short-distance nest burrow relocation to minimize construction impacts. *The Journal of Raptor Research.* 35(4):385-391 pp.
- Snyder, N.F.R. and J. Wiley. 1976. Sexual size dimorphism in hawks and owls of North America. *Ornithol. Monogr.* No. 20.
- Takats, D.L., G.L. Holroyd, J.R. Duncan, K.M. Mazur, C.M. Francis, R.J. Cannings, and W. Harris. 1999. Canadian nocturnal owl monitoring. *Proceedings of National Nocturnal owl Monitoring meeting in Winnipeg on September 27-28, 1999.* 16 pp.
- Thomsen, L. 1971. Behavior and ecology of Burrowing Owls on the Oakland Municipal Airport. *Condo.* 73:177-192.
- Trulio, L. 1995. Passive relocation: a method to preserve Burrowing Owls on disturbed sites. *J. Field Ornithol.* 66(1):99-106.

- Trulio, L. 1996. The western Burrowing Owl (*Speotyto cunicularia hypugaea*). BO Document Department of Geography and Environmental Studies, San Jose State University. 1-17 pp.
- Trulio, L. 1997. Burrowing Owl demography and habitat use at two urban sites in Santa Clara County, California. *J. Raptor Res. Report.* 9:84-89.
- Trulio, L. 1998. The Burrowing Owl as an indicator of CEQA effectiveness and environmental quality in the Silicon Valley. *In Environmental Monitor.* 4-5 pp.
- Trulio, L. and D. Rosenberg. 1998. Research on the demographic characteristics of Burrowing Owl populations in California: a progress report. *In Second international Burrowing Owl symposium.* The Raptor Research Foundation, Inc. 3 pp.
- U.S. Fish & Wildlife Service. 2001. Draft of Status assessment and conservation plan for the western Burrowing Owl in the United States. United States Department of the Interior, U.S. Fish & Wildlife. 145 pp.
- Walton, B.J. 1998. Burrowing Owl – Can the Endangered Species Act help? Presented at Second International Burrowing Owl Symposium. The Raptor Research Foundation, Inc. Ogden, UT. 5 pp.
- Wellicome, T.I. and G.L. Holroyd. 2001. The Second International Burrowing Owl Symposium: background and context. *J. Raptor Res.* 35(4): 269-273.
- Wellicome, T.I. and R.G. Poulin. 1998. Can we manage reproductive output in Burrowing Owls by managing their prey? *In Second international Burrowing Owl symposium.* The Raptor Research Foundation, Inc. 4 pp.
- Wellicome, T.I., G.L. Holroyd and H.E. Trefry. 1999. Are breeding populations of the Western Burrowing owl (*Athene cunicularis hypugaea*) declining throughout North America. *In Raptor Research Foundation Annual Meeting.* November 3-7, 1999, La Paz, Baja California Sur, Mexico. 13 pp.
- Winchell, C. S. 1994. Natural history and protection of burrowing owls. *Proceedings of the Vertebrate Pest Conference* 16: 83-86.
- Woollett, J.S. and M.G. van Hattem. 1999. Western burrowing owl demographic and biogeographic traits in a coastal range of central California. Presented at the Raptor Research Foundation Annual Meeting, 3-7 November 1999. La Paz, Baja, Mexico.
- WRI (Wildlife Research Institute, Inc.). 2003. Burrowing Owl Surveys on City of San Diego Properties (for the period January 1, 2001 – March 24, 2003). Prepared for City of San Diego. 24 March.

- WRI (Wildlife Research Institute, Inc.). 2005. Burrowing Owl Management and Monitoring Plan Lower Otay Lake Burrowing Owl Management Area. Prepared for City of San Diego. 31 March.
- WRI (Wildlife Research Institute, Inc.). 2002. Year 1 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2001). Prepared for California Department of Fish and Game. 30 July.
- WRI (Wildlife Research Institute, Inc.). 2004. Year 2 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2002). Prepared for California Department of Fish and Game. 19 March.
- WRI (Wildlife Research Institute, Inc.). 2005. Final Report – NCCP/MSCP Raptor Monitoring Project (2001-2003). Prepared for California Department of Fish and Game. 31 March.
- York, M.M., D.K. Rosenberg, and K.K. Sturm. 1997. Diet and food-niche breadth of Burrowing Owls (*Athene cunicularis*). U.S. Fish and Wildlife Service. 21 pp.
- Zarn, M. 1974. Burrowing Owl, Report No. 11. Habitat management series for unique or endangered species. Bureau of Land Management, U.S. Department of the Interior, Denver, CO. 25 pp.

Cooper's Hawk

- Anderson, D.W. and J.J. Hickey. 1972. Eggshell changes in certain North American birds. Proc. Int. Ornithol. Congr. 15:514-540.
- Boal, C.W. and R.W. Mannan. 1998. Nest site selection by Cooper's hawks in an urban environment. *J. Wildl. Manage.* 62:864-871.
- Brown, W.H. 1973. Winter population trends in Marsh, Cooper's and Sharp-shinned hawks. *Amer. Birds* 27:6-7.
- California Wildlife. 1982. Cooper's hawk *Accipiter cooperii*. *California's Wildlife*. 2:128-129.
- DeLong, J. and S.W. Hoffman. 1999. Differential autumn migration of Sharp-shinned and Cooper's Hawks in western North America. *Condor*. 101:674-678.
- Iverson, G.C. and M.R. Fuller 1992. Area-occupied survey technique for nesting woodland raptors. Proceedings of the Midwest Raptor Management Symposium and Workshop. Woodland Raptor Surveys Institute for Wildlife Research. National Wildlife Federation Scientific and Technical Series No. 15:1-7.
- Lincer, J.L. (Consulting Editor). 1989. Raptor Habitat Management Under the U.S. Bureau of Land Management Multiple-Use Mandate by R.R. Olendorff *et al.* Raptor Research Report No. 8, Raptor Research Foundation, Inc. Allen Press. 80 pp.

- Lincer, J.L. and R.J. Clark. 1978. Organochlorine residues in raptor eggs in the Cayuga Lake Basin, New York. *New York Fish Game J.* 25:121-128.
- Mosher, J.A. and M.R. Fuller. 1996. Surveying woodland hawks with broadcasts of great horned owl vocalizations. Reprinted from *Wildlife Society Bulletin*. 24(3):531-536.
- Mosher, J.A., M.R. Fuller, and M. Kopeny. 1990. Surveying woodland raptors by broadcast of conspecific vocalizations. *J. Field Ornithol.*, 61(4):453-461.
- Ogden (Ogden Environmental and Energy Services Co.). 1992. Otay Ranch Raptor Management Study. Submitted to Otay Ranch Project Team, Chula Vista, California.
- Remsen, Jr., J.V. 1978. Bird Species of Special Concern in California. California Department of Fish and Game, State of California, The Resources Agency, Department of Fish and Game. Wildlife Mgmt. Branch, Admin. Report No. 78-1. Pp.:31-32.
- Risebrough, R.W., R.W. Reiche, D.B. Peakall, S.G. Herman, and M.N. Kirven. 1968. Polychlorinated biphenyls in the global ecosystem. *Nature* 220:1098-1102.
- Rosenfield, R.N., J. Bielefeldt, and R.K. Anderson. 1988. Effectiveness of broadcast calls for detecting breeding Cooper's Hawks. *Wildl. Soc. Bull.* 16(2):210-212.
- Rosenfield, R.N. and J. Bielefeldt. 1993. Cooper's hawk *Accipiter cooperii*. . In *The Birds of North America*, No. 75 (A. Poole and F. Gill, eds.). The Academy of Natural Sciences, Philadelphia, PA and The American Ornithologists' Union, Washington, D.C. 24pp.
- Rosenfield, R.N., J. Bielefeldt, and R.K. Anderson. 2000. Comparative breeding ecology of Cooper's Hawk in urban vs. exurban areas of southeastern Arizona. *J. Wildl. Manage.* 64(2):599-600.
- Snyder, H.A and N.F.R. Snyder. 1974. Increased mortality of Cooper's hawks accustomed to man. *Condor*. 76:215-216.
- Snyder, N.F.R., H.A. Snyder, J.L. Lincer, and R.T. Reynolds. 1973. Organochlorines, heavy metals and the biology of North American accipiters. *BioScience*, 23(5):300-305.
- Walton, B.J., L.R. Mewaldt, and E.V. Johnson. 1976. Observations on Cooper's hawk (*Accipiter cooperii*) populations in California. 1972-1975. Unpubl. Rep. 12 pp.
- WRI (Wildlife Research Institute, Inc.). 2002. Year 1 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2001). Prepared for California Department of Fish and Game. 30 July.
- WRI (Wildlife Research Institute, Inc.). 2004. Year 2 Report for NCCP Raptor Monitoring

Project (January 1 – December 31, 2002). Prepared for California Department of Fish and Game. 19 March.

WRI (Wildlife Research Institute, Inc.). 2005. Final Report – NCCP/MSCP Raptor Monitoring Project (2001-2003). Prepared for California Department of Fish and Game. 31 March.

Ferruginous Hawk

Atkinson, E.C. 1995. Survey and monitoring guidelines for Ferruginous Hawks in Montana. Raptor Research Center, Boise State University, Boise, Id. Unpubl. Rep. 42 pp.

Ayers, L.W. and S.H. Anderson. 1999. An aerial sightability model for estimating Ferruginous Hawk population size. *J. Wildl. Manage.* 63(1):85-97.

Bechard, M.J. and J.K. Schmutz. 1995. Ferruginous Hawk *Buteo regalis*. The Birds of North America. 172:1-20.

CDFG. 1991. Ferruginous Hawk *Buteo regalis*. Species of Special Concern, California Dept. of Fish and Game. Unpubl. Rep. 13 pp

Garrison, B.A. 1990. Trends in winter abundance and distribution of Ferruginous Hawks in California. *Trans. West. Sect. Wildl. Soc.* 26:51-56.

Lincer, J.L. (Consulting Editor). 1989. Raptor Habitat Management Under the U.S. Bureau of Land Management Multiple-Use Mandate by R.R. Olendorff *et al.* Raptor Research Report No. 8, Raptor Research Foundation, Inc. Allen Press. 80 pp.

Lehman, R.N., L.B. Carpenter, K. Steenhof, and M.N. Kochert. 1991. Assessing relative abundance and reproductive success of shrubsteppe raptors. *J. Field Ornithol.* 69(2):244-256.

McAnnis, D.M. 1990. Home range, activity budgets, and habitat use of Ferruginous Hawks (*Buteo regalis*) breeding in southwest Idaho. MS Thesis, Boise State University, Boise 81pp.

Moritsch, M.Q. 1985. Photographic guide for aging nestling Ferruginous Hawks. Unpubl. Rep. U.S. Bureau of Land Management, Boise, Idaho. 22 Pp.

Ogden (Ogden Environmental and Energy Services Co.). 1992. Otay Ranch Raptor Management Study. Submitted to Otay Ranch Project Team, Chula Vista, California.

Olendorff, R.R. 1993. Status, biology, and management of Ferruginous Hawks; a review. Raptor Res. And Tech. Asst. Center, Spec. Rep. U.S Dept. Interior, Bur. Land Manage., Boise, Id. 84 pp.

Steenhof, K. 1984. Use of an interspecific communal roost by wintering Ferruginous Hawks. *Wilson Bull.* 96:137-138.

- Steenhof, K. and M.N. Kochert. 1985. Dietary shifts of sympatric buteos during a prey decline. *Oecologia* (Berlin) 66:6-16.
- Tate, J. 1986. The Blue List for 1986. *Am. Birds* 40:227-236.
- Tigner, J.R., M.W. Call and M.N. Kochert. 1996. Effectiveness of artificial nesting structures for Ferruginous Hawks in Wyoming. *In* Academic Press Ltd. 15:137-144
- Woffinden, N.D. and J.R. Murphy. 1989. Decline of a Ferruginous Hawk population: a 20-year summary. *J. Wildl. Manage.* 53(4):1127-1132.
- Woffinden, N.D. No date. A decade long extinction of a central Utah population of the Ferruginous Hawk *Buteo regalis*. Division of Natural Sciences, University of Pittsburgh, PA. Unpubl. Rep. 9 pp.
- WRI (Wildlife Research Institute, Inc.). 2002. Year 1 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2001). Prepared for California Department of Fish and Game. 30 July.
- WRI (Wildlife Research Institute, Inc.). 2004. Year 2 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2002). Prepared for California Department of Fish and Game. 19 March.
- WRI (Wildlife Research Institute, Inc.). 2005. Final Report – NCCP/MSCP Raptor Monitoring Project (2001-2003). Prepared for California Department of Fish and Game. 31 March.

Golden Eagle

- Anderson, D.W. and J.J. Hickey. 1972. Eggshell changes in certain North American birds. *Proc. Int. Ornithol. Congr.* 15:514-540.
- Bittner, J.D. 1997. Golden Eagles in San Diego County. Presented at the Raptor Research Foundation Annual Meeting. Ogden, UT.
- Bittner, J.D. and J. Oakley. 1999. Status of Golden Eagles in Southern California. Raptor Research Foundation Conference. Golden Eagle Symposium. November 2. La Paz, Baja California.
- Bittner, J.D., J.L. Lincer, J. Oakley, Nick Muscolino, and J. Hannan. 2003. Golden Eagle reproduction in a drought period. Presented at Raptor Research Foundation's Annual Scientific Conference. September 2-7. Anchorage, AK.
- Bloom, P. H. 1991. Status of the Golden Eagle population on Marine Corps Base Camp Pendleton. Unpubl. Rep. Santa Ana, California. 21 pp.

- California. Department of Fish and Game. 1982. Golden Eagle *Aquila chrysaetos*. California's Wildlife. 2:142 p.
- Collopy, M.W. 1983. A comparison of direct observations and collections of prey remains in determining the diet of Golden Eagles. *J. Wildl. Manage.* 47:360-368.
- Collopy, M.W. 1983. Foraging behavior and success of Golden Eagles. *Auk* 100:747-749.
- Collopy, M.W. 1986. Food consumption and growth energetics of nestling Golden Eagles. *Wilson Bull.* 98:445-458.
- Collopy, M.W., and T.C. Edwards Jr. 1989. Territory size, activity budget, and role of undulating flight in nesting Golden Eagles. *J. Field Ornithol.* 60:43-51.
- DeSmet, K.D. 1987. Status report on the Golden Eagle *Aquila chrysaetos*. Committee on the status of endangered wildlife in Canada, Ottawa, ON, Status assigned in 1982, reviewed in 1995. Unpubl. Rep.
- Dixon, J.B. 1937. The Golden Eagle in San Diego County, California *The Condor.* 39(2)49-56.
- Edwards, T.C., Jr., M.W. Collopy, K. Steenhof, and M.N. Kochert. 1988. Sex ratios of fledgling Golden Eagles. *Auk* 105:793-796.
- Edwards, T.C., Jr., and M. N. Kochert. 1986. Use of body weight and length of footpad as predictors of sex in Golden Eagles. *J. Field Ornithol.* 57:317-319.
- Harlow, D.L. and P.H. Bloom. 1989. Buteos and the Golden Eagle. *In* Western Raptor Management Symposium and Workshop. 102-110 pp.
- Harmata, A.R. 1993. Heavy metal and pesticide contamination of Bald and Golden Eagles in the western United States. Unpubl. USEPA, December 1993. 43 pp.
- Hinds, K.O. and H. de la Cueva. 1999. Habitat and competitive abilities of the Golden Eagle (*Aquila chrysaetos*), in Sierra San Pedro Martir, Baja California. *In* Raptor Research Foundation Annual Meeting, November 3-7. La Paz, Baja California Sur, Mexico. 20 pp.
- Hoechlin, D.R. 1974. Behavioral ecology of nesting Golden Eagles (*Aquila chrysaetos*) in San Diego County. M.S. Thesis, San Diego State University, San Diego, California 113 pp.
- Hunsicker, G.R. 1972. Nesting Behavior of the Golden Eagle, *Aeuila chrysaetos*, in San Diego County, California. M.S. Thesis, University of California Riverside 65 pp.
- Kirk, D.A. 1996. Updated status report on the Golden Eagle *Aeuila chrysaetos*. Committee on the Status of Endangered Wildlife in Canada, Ottawa, Ontario, Canada, Status assigned in 1996.

- Kochert, M. N. 1980. Golden Eagle reproduction and population changes in relation to jackrabbit cycles: Implications to eagle electrocutions. pp. 71-86. *In* R. P. Howard and J. F. Gore [eds.], A workshop on raptors and energy developments. U.S. Fish and Wildlife Service and the Idaho Chapter of the Wildlife Society, Boise, ID. (RRTAC Reprint #2)
- Kochert, M.N. and K. Steenhof. 1999. Golden Eagles in the U.S. and Canada; status, trends conservation challenges. *Raptor Res. Rep.* 11:1-19. (In Review: Raptor Research Report No. 11. Proceedings of the Golden Eagle Symposium – La Paz, Mexico, 2 Nov. 99).
- Kochert, M.N., K. Steenhof, L.B. Carpenter, and J.M. Marzluff. 1999. Effects of fire on Golden Eagle territory occupancy and reproductive success. *J. Wildl. Manage.* 63:773-780.
- Kochert, M.N., K. Steenhof, C.L. McIntyre, and E.H. Craig. 2002. Golden Eagle (*Aquila chrysaetos*). *In* The Birds of North America, No. 684 (A. Poole and F. Gill, eds.). The Academy of Natural Sciences, Philadelphia, PA and The American Ornithologists' Union, Washington, D.C.
- Lederle, P.E., J.M. Mueller, and E.A. Holt. 2000. Raptor surveys in south central Nevada, 1991-95. *J. Raptor Res.* 34(2): 133-136.
- Lincer, J.L. (Consulting Editor). 1989. Raptor Habitat Management Under the U.S. Bureau of Land Management Multiple-Use Mandate by R.R. Olendorff *et al.* Raptor Research Report No. 8, Raptor Research Foundation, Inc. Allen Press. 80 pp.
- Lincer, J.L. and J.D. Bittner. 2002. Use of Raptors to Monitor HCPs. Poster presented at the AOU/Raptor Research Foundation, New Orleans, September 24-28.
- Lincer, J.L., J.D. Bittner, N. Muscolino, and, L. Swartz. 2003. A Raptor Protocol for Monitoring HCPs. Paper presented at the Raptor Research Foundation's Annual Scientific Conference. September 2-7. Anchorage, AK.
- Lockhart, J.M. and M.N. Kochert. 1978. Effects of visual markers and telemetry devices on the nesting success of Golden Eagles. Draft manuscript. 14 pp.
- Marzluff, J.M., S.T. Knick, M.S. Vekasy, L.S. Schueck, and T.J. Zarriello. 1997. Spatial use and habitat selection of Golden Eagles in southwestern Idaho. *Auk* 114:673-687.
- Marzluff, J.M., M.S. Vekasy, M.N. Kochert, and K. Steenhof. 1997. Productivity of Golden Eagles wearing backpack radiotransmitters. *J. Raptor Res.* 31:223-227.
- Mosher, J.A. and C.M. White. 1976. Directional exposure of Golden Eagles nests. *Can. Field-Nat.* 90:356-359.

- Murphy, J.R. 1977. Eagles and livestock-Some management considerations. pp. 307-314. *In* World Conference on Birds of Prey. Vienna, 1-3 October, 1975. Report of proceedings, (R.D. Chancellor, ed.) International Council for Bird Preservation, London.
- Ogden (Ogden Environmental and Energy Services Co.). 1992. Otay Ranch Raptor Management Study. Submitted to Otay Ranch Project Team, Chula Vista, California.
- Olendorff, R.R. 1975. Golden Eagle county, 1st ed. Alfred A. Knopf, Inc., New York.
- Opdycke, J.D. 1993. Potential impacts to Golden Eagles on Iron Mountain, Ramona, San Diego County, California. Fish and Wildlife Service, Ecological Services. Unpubl. Rep. 3 pp.
- Phillips, R.L., J.L. Cummings, and J.D. Berry. 1991. Responses of breeding Golden Eagles to relocation. *Wildl. Soc. Bull.* 19:430-434.
- Remsen, J. V. 1978. *Bird species of special concern in California: an annotated list of declining or vulnerable bird species*. Nongame Wildlife Investigations, Wildlife Management Branch, California Department of Fish & Game. Administrative Report No. 78-1.
- Remsen, V. 1980. *Bird Species of Special Concern in California*. California Department of Fish and Game, Sacramento, California, 54 pp.
- Schlorff, R.W. 1986. Nongame Wildlife Investigations, Golden Eagle status review. Job Final Report. State of California, The Resource Agency, Department of Fish and Game. W-65-R-2:11.
- Schueck, L.S., J.M. Marzluff, M. Vekasy, M.R. Fuller, T.J. Zarriello, and W.S. Seegar. 1995. Abstract: Migration routes and winter ranges of Golden Eagles. *J. Raptor Res.* 29:72-73.
- Scott, T.A. 1982. Human impacts on the Golden Eagle population of San Diego County from 1928 to 1981. M.S. Thesis, San Diego State University, San Diego, California 101 pp.
- Small, A. 1976. Development of Golden Eaglets in southern California. *Western Birds.* 7:137-152.
- Snow, C. 1973. Habitat management series of unique or endangered species. Report No. 7. Bureau of Land Management, Denver, Colorado. 52 pp.
- Spofford, W.R. 1969. Brief reports: The status of eagles. Problems of the Golden Eagles in North America. pp. 345-347. *In* Peregrine Falcon populations: their biology and decline, (J.J. Hickey, ed.). University of Wisconsin Press, Madison.
- Steenhof, K., M.N. Kochert, and J.H. Doremus. 1983. Nesting of subadult Golden Eagles in southwestern Idaho. *Auk* 100:743-747.

- Steenhof, K., M.N. Kochert, and T.L. McDonald. 1997. Interactive effects of prey and weather on Golden Eagle reproduction. *J. Anim. Ecol.* 66:350-362.
- Thelander, C.G. 1974. Nesting territory utilization by Golden Eagles (*Aquila chrysaetos*) in interior Central Coast Ranges of California. Jones & Stokes Associates, Inc., Sacramento, CA.
- Thelander, C.G. 1974. Nesting territory utilization by golden eagles in California during 1974. California Department of Fish and Game, Wildl. Manage. Branch, Admin. Report No. 74-7. 19pp.
- Watson, J. 1997. The Golden Eagle. T & AD Poyser. London. 374 pp.
- WRI (Wildlife Research Institute, Inc.). 2002. Year 1 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2001). Prepared for California Department of Fish and Game. 30 July.
- WRI (Wildlife Research Institute, Inc.). 2004. Year 2 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2002). Prepared for California Department of Fish and Game. 19 March.
- WRI (Wildlife Research Institute, Inc.). 2005. Final Report – NCCP/MSCP Raptor Monitoring Project (2001-2003). Prepared for California Department of Fish and Game. 31 March.
- WRI (Wildlife Research Institute, Inc.). 2005. Final Report for NCCP/MSCP Raptor Monitoring Project (2001-2003); Golden Eagles of the San Diego MSCP Area. Prepared for California Department of Fish and Game. 31 March.

Northern Harrier

- California. Department of Fish and Game. 1982. Northern Harrier *Circus cyaneus*. California's Wildlife. 2:124 p.
- California. Department of Fish and Game. 1987. Job Final Report. Nongame Wildlife Investigations, Northern Harrier Breeding Survey, Job No. II-18. 6pp.
- California Department of Fish and Game. 1990. California statewide wildlife habitat relationships system. Volume II: Birds, D. Zeiner, W. Laudenslayer, K. Mayer, and M. White (eds.). The Resource Agency, Sacramento. 731 pp.
- Collopy, M.W. and K.L. Bildstein. 1987. Foraging behavior of Northern Harriers wintering in southeastern salt and freshwater marshes. *Auk*. 104:11-16.
- Estep, J.A. 1986. Nongame Wildlife Investigations, Marsh hawk status survey. Job Progress Report. State of California, The Resource Agency, Department of Fish and Game. W-65-R-3:5.

- Larsen, C.J. 1987. Nongame Wildlife Investigations, Northern Harrier breeding survey. Job Final Report. State of California, The Resource Agency, Department of Fish and Game. W-65-R-4:6.
- Lederle, P.E., J.M. Mueller, and E.A. Holt. 2000. Raptor surveys in southcentral Nevada, 1991-95. *J. Raptor Res.* 34(2):133-136.
- Lehman, R.N., L.B. Carpenter, K. Steenhof, and M.N. Kochert. 1991. Assessing relative abundance and reproductive success of shrubsteppe raptors. *J. Field Ornithol.* 69(2) 244-256.
- Lincer, J.L. (Consulting Editor). 1989. Raptor Habitat Management Under the U.S. Bureau of Land Management Multiple-Use Mandate by R.R. Olendorff *et al.* Raptor Research Report No. 8, Raptor Research Foundation, Inc. Allen Press. 80 pp.
- Macwhirter, R.B. and K.L. Bildstein. 1996. Northern Harrier *Circus cyaneus*. In *The Birds of North America*, No. 210 (A. Poole and F. Gill, eds.). The Academy of Natural Sciences, Philadelphia, PA and The American Ornithologists' Union, Washington, D.C.
- Martin, J.W. 1987. Behavior and habitat use of breeding Northern Harriers in southwestern Idaho. *J. Raptor Res.* 21(2):57-66.
- Nesbitt, S.A. 1985. Northern Harriers. *Eyas* 8(2) 28-29.
- Ogden (Ogden Environmental and Energy Services Co.). 1992. Otay Ranch Raptor Management Study. Submitted to Otay Ranch Project Team, Chula Vista, California.
- Preston, C.R. 1990. Distribution of raptor foraging in relation to prey biomass and habitat structure. *Condor.* 92:107-122.
- Remsen, Jr., J.V. 1978. Bird Species of Special Concern in California. California Department of Fish and Game, Wildlife Mgmt. Branch, Admin. Report No. 78-1. 54 pp.
- WRI (Wildlife Research Institute, Inc.). 2002. Year 1 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2001). Prepared for California Department of Fish and Game. 30 July.
- WRI (Wildlife Research Institute, Inc.). 2004. Year 2 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2002). Prepared for California Department of Fish and Game. 19 March.
- WRI (Wildlife Research Institute, Inc.). 2005. Final Report – NCCP/MSCP Raptor Monitoring Project (2001-2003). Prepared for California Department of Fish and Game. 31 March.

Peregrine Falcon

- Anderson, D.W. and J.J. Hickey. 1972. Eggshell changes in certain North American birds. *Proc. Int. Ornithol. Congr.* 15:514-540.
- Banks, R.C. 1969. Peregrine Falcon in Baja California and the Gulf of California, Peregrine Falcon populations: their biology and decline, (J. J. Hickey, ed.). University of Wisconsin Press, Madison. 81-91 pp.
- Cade, T.J., J.L. Lincer, C.M. White, D.G. Roseneau and L.G. Swartz. 1971. DDE residues and eggshell changes in Alaskan falcons and hawks. *Science*, 172:955-957.
- Cade, T, J. Enderson, C. Thelander, and C. White (Eds.). 1987. Peregrine Falcon Populations: Their Management and Recovery. Published by the Peregrine Fund, Inc.
- California. Department of Fish and Game. 1980. American Peregrine Falcon *Falco peregrinus anatum*. *At the crossroads*. 91-92 pp.
- California. Department of Fish and Game. 1982. Peregrine Falcon *Falco peregrinus*. California's Wildlife. 2:148 p.
- Dawson, W.L. 1923. The Peregrine Falcon (*Falco peregrinus anatum*). Agency Review Draft. 24 pp.
- Fuller, M.R., W.S. Seegar, and L.S. Schueck. 1998. Routes and travel rates of migrating Peregrine Falcons *Falco peregrinus* and Swainson's Hawks *Buteo swainsoni* in the western hemisphere. *Journal of Avian Biology* 29:433-440. (Proceedings: "Optimal Migration", Lund University, Lund, Sweden, November 5-8, 1997).
- Fyfe, R., R.W. Risebrough, J.G. Monk, W.M. Jarman, D.W. Anderson, L.F. Kiff, J.L. Lincer, I.C.T. Nisbet, W. Walker II, and B.J. Walton. 1987. DDE, Productivity and Eggshell Thickness Relationships Within the Genus *Falco*. Chapter 33 *In* Peregrine Falcon Populations: Their Management and Recovery. Edited by T. Cade, J. Enderson, C. Thelander, C. White. Published by the Peregrine Fund, Inc.
- Hays, L. and S. Grandberry. 1998. The Peregrine Falcon is back: babbitt announces proposal to remove world's fastest bird from endangered species list. U.S. Fish and Wildlife Service, Carlsbad, California.
- Hickey, J. J. (ed.). 1969. Peregrine Falcon populations: their biology and decline, University of Wisconsin Press, Madison.
- Jurek, R.M. 1989. Five year status report-American Peregrine Falcon. State of California, The Resource Agency, Department of Fish and Game. 1-15 pp.
- Lincer, J.L. 1975. The effects of dietary DDE on eggshell-thinning in the American kestrel: A comparison of the field situation and laboratory results. *J. Applied Ecol.*, 12(3): 781-793.

- Lincer, J.L. (Consulting Editor). 1989. Raptor Habitat Management Under the U.S. Bureau of Land Management Multiple-Use Mandate by R.R. Olendorff *et al.* Raptor Research Report No. 8, Raptor Research Foundation, Inc. Allen Press. 80 pp.
- McWilliams, S.R., J.P. Dunn, and D.G. Raveling. 1994. Predator-prey interactions between eagles and cackling Canada and Ross' geese during winter in California. *Wilson Bull.* 106:272-288.
- Porter, R.D., A.M. Jenkins, M.N. Kirven, D.W. Anderson, and J.O. Keith. 1988. Status and reproductive performance of marine peregrines in Baja California and the Gulf of California, Mexico. *In* Peregrine Falcon populations: Their management and recovery, (T. J. Cade, J. H. Enderson, C. G. Thelander, and C. M. White, eds.). The Peregrine Fund, Inc., Boise, ID. 105-114 pp.
- Smith, B. 1989. Plan for Bald Eagles sought at Cachuma. Santa Barbara News Press (Santa Barbara, CA) (March 6): A1, A5.
- Steinhof, P. 1990. American Peregrine Falcon *Falco peregrinus anatum*. California Wildl. Heritage. Threatened and endangered animals in the golden state. Department of Fish and Game. 27-28.
- Thelander, C.G. 1975. Distribution and reproductive success of Peregrine Falcons (*Falco peregrinus anatum*) in California-1975. State of Calif. Resour. Agency. Dept. Fish and Game. Wildl. Manag. Branch. Admin. Rep. No. 75-6. 12 pp.
- Thelander, C.G. 1976. Distribution and reproductive success of Peregrine Falcons (*Falco peregrinus anatum*) in California during 1975 and 1976. State of Calif. Resour. Agency. Dept. Fish and Game. Wildl. Manag. Branch. Admin. Rep. No. 76-3. 13 pp.
- Thelander, C.G. 1977. The breeding status of Peregrine Falcons in California. MA Thesis, San Jose State Univ., San Jose, California 112 pp.
- Walton, B.J. 1981. Peregrine Falcon management in California - update for 1981. Unpubl. rep., Santa Cruz Predatory Bird Res. Group, Univ. of California, Santa Cruz. 58 pp.
- Watson, J.F. 1981. Ecological characterization of the central and northern California coastal region. Volume II, Part 2. Species. Unpub. rep. FWS/OBS/46.2, October 1981. Prepared by Jones & Stokes Assoc., Inc., Sacramento, California. (Pages II-151 and II-152 only)
- Wendt, A., G. Septon, and J. Moline. 1991. Juvenile urban-hacked Peregrine Falcons (*Falco peregrinus*). *J. Raptor Res.* 25:94-95.
- Wootton, J.T., and D.A. Bell. 1992. A metapopulation model of the Peregrine Falcon in California: viability and management strategies. *Ecol. Appl.* 2(3):307-321.

- WRI (Wildlife Research Institute, Inc.). 2002. Year 1 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2001). Prepared for California Department of Fish and Game. 30 July.
- WRI (Wildlife Research Institute, Inc.). 2004. Year 2 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2002). Prepared for California Department of Fish and Game. 19 March.
- WRI (Wildlife Research Institute, Inc.). 2005. Final Report – NCCP/MSCP Raptor Monitoring Project (2001-2003). Prepared for California Department of Fish and Game. 31 March.

Swainson's Hawk

- Bloom, P.H. 1980. The status of the Swainson's Hawk in California. State of California, The Resource Agency, Department of Fish and Game. Bureau of Land Management, and Federal Aid in Wildlife Restoration, Project W-54-R-12. 42 pp.
- California Department of Fish and Game. 1982. Swainson's Hawks *Buteo swainsoni*. California's Wildlife. 2:134 p.
- California Department of Fish and Game 1988. Five year status report Swainson's Hawks *Buteo swainsoni*. Nongame Bird and Mammal Section, Wildlife Management Division, 1-9 pp.
- California Department of Fish and Game. 1990. Mitigation criteria for Swainson's Hawks, Region 2, State of California. 2 pp.
- California Department of Fish and Game. 1990. Mitigation guidelines for Swainson's Hawks *Buteo swainsoni* in the Central Valley of California. Region 2, State of California. 1-12 pp.
- Cox, J.D. 1997. Surprised researchers find California hawks don't join in migration to Argentina. The San Diego Union-Tribune. Scripps-McClatchy Western Service.
- Estep, J.A. 1989. Biology, movements, and habitat relationships of the Swainson's Hawk in the Central Valley of California, 1986-87. Wildlife Management Division, Nongame Bird and Mammal Section, State of California, The Resource Agency, Department of Fish and Game. 51 pp.
- England, A.S., M.J. Bechard, and C. S. Houston. 1997. Swainson's Hawks *Buteo swainsoni*. In The Birds of North America, No. 265 (A. Poole and F. Gill, eds.). The Academy of Natural Sciences, Philadelphia, PA and The American Ornithologists' Union, Washington, D.C. 27 pp.

- England, A.S., J.A. Estep, and W.R. Holt. 1995. Nest-site selection and reproductive performance of urban-nesting Swainson's Hawks in the Central Valley of California. *J. Raptor Res.* 29:179-186.
- Fuller, M.R., W.S. Seegar, and L.S. Schueck. 1998. Routes and travel rates of migrating Peregrine Falcons *Falco peregrinus* and Swainson's Hawks *Buteo swainsoni* in the western hemisphere. *Journal of Avian Biology* 29:433-440. (Proceedings: "Optimal Migration", Lund University, Lund, Sweden, November 5-8, 1997).
- Goldstein, D.L., and N.G. Smith. 1991. Short communications. *J. Raptor Res.* 25(3):87-88.
- Hall, R.S. No date. Preliminary status report and notes on the breeding biology of the Swainson's Hawk in Northwest Arizona. *In Press.* 1-6 pp.
- Lincer, J.L. (Consulting Editor). 1989. Raptor Habitat Management Under the U.S. Bureau of Land Management Multiple-Use Mandate by R.R. Olendorff *et al.* Raptor Research Report No. 8, Raptor Research Foundation, Inc. Allen Press. 80 pp.
- Risebrough, R.W., R.W. Schlorff, P.H. Bloom, and E.E. Littrell. 1989. Investigations of the decline of Swainson's Hawk populations in California. *J. Raptor Res.* 23(3):63-71.
- Schlorff, R.W. 1985. Diurnal raptor population monitoring program. Nongame Wildlife Program, State of California, Department of Fish and Game. W-65-R-2:24 pp.
- Schlorff, R.W. and P.H. Bloom. 1980. Importance of Riparian systems to nesting Swainson's Hawks in the Central Valley of California. Nongame Wildlife Program, State of California, Department of Fish and Game. 13 pp.
- Sharp, B. 1986. Management guidelines for the Swainson's Hawk. Region 1. U.S. Fish and Wildlife Service, Portland, Oregon. 1-28 pp.
- Sharp, C.S. 1902. Nesting of Swainson Hawk. *Condor* 4:116-118.
- Woodbridge, B. 1991. Habitat selection by nesting Swainson's Hawks: a hierarchical approach. M.S. Thesis. Oregon State University. 80 pp.
- WRI (Wildlife Research Institute, Inc.). 2002. Year 1 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2001). Prepared for California Department of Fish and Game. 30 July.
- WRI (Wildlife Research Institute, Inc.). 2004. Year 2 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2002). Prepared for California Department of Fish and Game. 19 March.
- WRI (Wildlife Research Institute, Inc.). 2005. Final Report – NCCP/MSCP Raptor Monitoring Project (2001-2003). Prepared for California Department of Fish and Game. 31 March.

General and Raptor Management Literature

- Anderson, D.E., G.J. Hongstad, and W.R. Myiton. 1985. Line transect analysis of raptor abundance along roads. *Wildl. Soc. Bull.* 13:533-539.
- Avian Power Line Interaction Committee (APLIC). 1994. Mitigating bird collisions with power lines: the state of the art in 1994. Edison Electric Institute. Washington, D.C. 78pp.
- Avian Power Line Interaction Committee (APLIC). 1996. Suggested Practices for Raptor protection on power Lines: State of the Art in 1996. Edison Electric Institute/Raptor Research Foundation. Washington, D.C.
- Barrows, C.W. 1989. Diets of five species of desert owls. *Western Birds.* 20:1-10.
- Bloom, P.H. 1985. Raptor movements in California. In M. Harwood [Ed.], Proceedings of Hawk Migration Association of North America, North Wales, PA U.S.A. 313-324 pp.
- Buckland, K.P., D.R. Anderson and J.L. Laake. 1981. Line transect estimation of bird population density using a Fourier series. *Stud. Avian Biol.* 6:466-482.
- Buckland, S.T. 1987. On the variable circular plot method of estimating animal density. *Biometrics* 43:363-384.
- Buckland, S.T., D.R. Anderson, K.P. Burnham, and J.L. Laake. 1993. Distance sampling: estimating abundance of biological populations. Chapman and Hall, London, U.K.
- Clark, W.S. and B.K. Wheeler. 1987. A Field Guide to Hawks North America. Houghton Mifflin Company. Boston.
- Everett, W.T. 1979. Threatened, declining and sensitive bird species in San Diego County. *Sketches* June.
- Fuller, M.R. and J.A. Mosher. 1981. Methods of detecting and counting raptors: a review. *Stud. Avian Biol.* 6:235-246.
- Gould, W. R., and M. R. Fuller. 1995. Survival and population size estimation in raptor studies: A comparison of two methods. *J. Raptor Res.* 29:256-264.
- Iverson, G.C. and M.R. Fuller 1992. Area-occupied survey technique for nesting woodland raptors. Proceedings of the Midwest Raptor Management Symposium and Workshop. Woodland Rapt. Surv. Institute for Wildlife Research. National Wildlife Federation Scientific and Technical Series No. 15:1-7.

- Kochert, M. N., K. Steenhof, and M. Q. Moritsch. 1983. Evaluation of patagial markers for raptors and ravens. *Wildl. Soc. Bull.* 11:271-281.
- Johnsgard, P.A. 1990. Hawks, eagles & falcons of North America: biology and natural history. Smithsonian Institution Press, Washington , D.C.
- Lehman, R. N., L. B. Carpenter, K. Steenhof, and M. N. Kochert. 1998. Assessing relative abundance and reproductive success of shrubsteppe raptors. *J. Field Ornithol.* 69:244-256.
- Lincer, J.L. 1982. "Protecting Endangered Species at the Local Governmental Level." Paper presented at the 46th Annual Meeting of the Florida Academy of Sciences, April 22-24, DeLand, FL. *Florida Scientist*, 45(1): 40.
- Lincer, J.L. 1983. "But Release Them to Where?" Raptor Research & Rehabilitation Program Newsletter, 4-(Winter). 1982-83:6-8.
- Lincer, J.L. 1984. The Priority of Proper Habitat Management. The EYAS (a newsletter of the National Wildlife Federation's Raptor Information Center), Vol. 7.
- Lincer, J.L. (Consulting Editor). 1989. Raptor Habitat Management Under the U.S. Bureau of Land Management Multiple-Use Mandate by R.R. Olendorff *et al.* Raptor Research Report No. 8, Raptor Research Foundation, Inc. Allen Press. 80 pp.
- McDermott, F. 1999. Pacific continental flyway. *HMANA Hawk Migration Stud.* 25:24-32.
- Mosher, J. A., and M. R. Fuller. 1996. Surveying woodland hawks with broadcasts of great horned owl vocalizations. *Wildl. Soc. Bull.* 24:531-536.
- Myers, R.I.G. Morrison, P.Z. Antas, B.A. Harrington, T.E. Lovejoy, M. Sallaberry, S.E. Senner, and A. Tarrak. 1987. Conservation strategy for migratory species. *Am. Sci.* 75:18.
- National Wildlife Federation. 1989. Proceedings of the western raptor management symposium and workshop. National Wildlife Federation, Washington, D.C., 320 pp.
- Newton, I. 1979. Population ecology of raptors. Buteo Books, Vermillion, SD USA.
- Ogden (Ogden Environmental and Energy Services Co.). 1992. Otay Ranch Raptor Management Study. Submitted to Otay Ranch Project Team, Chula Vista, California.
- Olendorff, R.R., A.D. Miller, and R. N. Lehman. 1981. Suggested practices for raptor protection on power lines--the state-of-the-art in 1981. Raptor Res. Rep. No. 4. Raptor Research Foundation, Inc. St. Paul Minn. 111pp.
- Reynolds, R.T., J.M. Scott, and R.A. Nussbaum. 1990. A variable circular-plot method for estimating bird numbers. *Condor* 82:309-313.

- Schueck, L. S., M. R. Fuller, and W. S. Seegar. 1989. Falcons. pp. 71-80. *In* B. G. Pendleton, M. N. LeFranc Jr., M. B. Moss, C. E. Ruibal, M. A. Knighton, and D. L. Krahe [eds.], Proceedings of the northeast raptor management symposium and workshop. National Wildlife Federation, Washington, DC. May 16-16, 1988, Syracuse, N.Y. (Scientific and Technical Series; no.13)
- Steenhof, K., and M.N. Kochert. 1982. An evaluation of methods used to estimate raptor nesting success. *J. Wildl. Manage.* 46:885-893.
- Steenhof, K., and M.N. Kochert. 1985. Dietary shifts of sympatric buteos during a prey decline. *Oecologia* 66:6-16.
- Steenhof, K., and M.N. Kochert. 1988. Dietary responses of three raptor species to changing prey densities in a natural environment. *J. Anim. Ecol.* 57:37-48.
- Steenhof, K., M.N. Kochert, and J.A. Roppe. 1993. Nesting by raptors and common ravens on electrical transmission line towers. *J. Wildl. Manage.* 57:271-281.
- Sutherland, W.J. 1996. Ecological census techniques. A handbook. Cambridge Univ. Press, Cambridge, U.K.
- Tate, J., Jr. and D.J. Tate. 1982. The Blue List for 1982. *Am. Birds* 36:126-135.
- Tate, J. 1986. The Blue List for 1986. *Am. Birds* 40:227-236.
- Williams, R. D. and E.W. Colson. 1989. Raptor associations with linear rights-of-way. Pages 173-192 *in* B. G. Pendleton, ed. Proc. Western Raptor Management Symp. Natl. wildl. Fed. Scientific and Tech. Series No. 12. Washington, D.C.
- Thiollay, J.-M. 1976. Les decompptes de rapaces le long des routes: essai de standardization. *Passer* 13:69-76.
- Unitt, P. 1984. The Birds of San Diego County. San Diego Society of Natural History, Memoir 13. Luster Industries.
- U.S. Department of the Interior. 1996. Effects of military training and fire in the Snake River Birds of Prey National Conservation Area. U.S. Geol. Survey, Biol. Res. Div., Snake River Field Station, Boise, ID, BLM/IDARNG Research Project Final Report.
- WRI (Wildlife Research Institute, Inc.). 2002. Year 1 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2001). Prepared for California Department of Fish and Game. 30 July.
- WRI (Wildlife Research Institute, Inc.). 2004. Year 2 Report for NCCP Raptor Monitoring Project (January 1 – December 31, 2002). Prepared for California Department of Fish and Game. 19 March.

WRI (Wildlife Research Institute, Inc.). 2005. Final Report for NCCP Raptor Monitoring Project (2001-2003). Prepared for California Department of Fish and Game. 31 March.

APPENDIX B

BREEDING SEASON RAPTOR NESTS AND TERRITORIES BY SITE (2001-2003)

The following pages reflect raptor breeding territories which were typical of the below study sites for the period 2001-2003. To facilitate the reader's access to the following topographic maps, they are listed below alphabetically and by site number.

<u>Number</u>	<u>Name</u>	<u>Name</u>	<u>Number</u>
1	Crestridge	Boden Canyon	2
2	Boden Canyon	Border Fields	44
3	Jamul Ranch	Brown Field Complex	22
4	SDNWR*/Salt Works/Egger Ghio	Crestridge	1
5	McGinty Mountain Complex	Grasslands/Route 67	30
6	San Diego Bay NWR (winter only)	Hollenbeck Canyon	34
7	Lake Hodges	Immenschuh	39
8	Penasquitos Lagoon	Iron Mountain	11
9	Torrey Pines	Jamul Ranch	3
10	Sycamore Canyon	Lake Hodges	7
11	Iron Mountain	Los Montanas (North)	40
12	Otay Mountain	Los Montanas (South)	41
13	Marron Valley	Marron Valley	13
14	Otay Lakes	McGinty Mountain Complex	5
15	SDNWR* Sweetwater Marsh	Miramar Reservoir	20
16	San Vicente	Mission Bay	21
17	Sycuan Peak	Mission Trails	24
18	Point Loma	North Island	19
19	North Island	Otay Lakes	14
20	Miramar Reservoir	Otay Mountain	12
21	Mission Bay	Penasquitos Canyon	33
22	Brown Field Complex	Penasquitos Lagoon	8
23	SDNWR*/San Miguel Mountain	Point Loma	18
24	Mission Trails	Proctor Valley	25
25	Proctor Valley	Rancho San Diego (East)	42
26	San Diego River	Rancho San Diego (West)	43
27	Route 67 South	Rock Mountain	35
28	San Dieguito Lagoon	Rockwood Canyon	32
29	Route S-6 (deleted/safety issue)	Route 67 South	27
30	Grasslands/Route 67	Route 94 (North and South)	38
31	Sloan Canyon	Route S-6 (deleted/safety issue)	29
32	Rockwood Canyon	San Diego Bay NWR (winter only)	6
33	Penasquitos Canyon	San Diego River	26
34	Hollenbeck Canyon	San Dieguito Lagoon	28
35	Rock Mountain	San Pasqual	36
36	San Pasqual	San Vicente	16
37	SDNWR*Tijuana Slough	SDNWR* Sweetwater Marsh	15
38	Route 94 (North and South)	SDNWR*/Salt Works/Egger Ghio	4
39	Immenschuh	SDNWR*/San Miguel Mountain	23
40	Los Montanas (North)	SDNWR*Tijuana Slough	37
41	Los Montanas (South)	Sloan Canyon	31
42	Rancho San Diego (East)	Sweetwater Reservoir	45
43	Rancho San Diego (West)	Sycamore Canyon	10
44	Border Fields	Sycuan Peak	17
45	Sweetwater Reservoir	Torrey Pines	9

*San Diego National Wildlife Refuge

LEGEND

Symbols

Center of raptor/corvid territory or assumed or documented nest site.

Note: Above symbol without an acronym following it indicates that a stick nest was documented but species was not determinable. If species was known for the nest or territory, the above symbol is followed by the appropriate acronym (see below).

Acronyms for Raptor and Corvid Species

AC	American crow
AK	American kestrel
BE*	BALD EAGLE
BH	Black hawk
BR	Barn owl
BO*	BURROWING OWL
CH*	COOPER'S HAWK
CR	Common raven
FH*	FERRUGINOUS HAWK
GE*	GOLDEN EAGLE
GO	Great-horned owl
HH	Harris' hawk
LO	Long-eared owl
MR	Merlin
NH*	NORTHERN HARRIER
OS	Osprey
PF*	PEREGRINE FALCON
PR	Prairie falcon
RS	Red-shouldered hawk
RT	Red-tailed hawk
SE	Short-eared owl
SO	Screech owl
SS	Sharp-shinned hawk
SH*	SWAINSON'S HAWK
TV	Turkey vulture
UA	Unidentifiable accipiter
UB	Unidentifiable buteo
UF	Unidentifiable falcon
UR	Unidentifiable raptor
WK	White-tailed kite
WH	White-tailed hawk
ZH	Zone-tailed hawk

* MSCP target species.

APPENDIX C

LONG -TERM RAPTOR MONITORING PROTOCOL

BACKGROUND

The Multiple Species Conservation Program (MSCP) is a comprehensive, long-term habitat conservation plan that addresses the needs of multiple species and the preservation of natural vegetation in San Diego County (County of San Diego 1997). The size and configuration of the preserve network is continually evolving but it may ultimately encompass approximately 172,000 acres. In order to determine if the MSCP or any management area, for that matter, is functioning correctly, a meaningful monitoring plan must be in place. A vast area, such as the MSCP, cannot be comprehensively monitored for any but a few species with very limited and specific habitat requirements. Raptor species will, therefore, be monitored using a reproducible sampling approach. Details of this approach are described below after reminding the reader of the ultimate monitoring objectives.

OBJECTIVES

The overall goal of the MSCP monitoring is to detect changes in habitat quality and population trends in those habitats and species covered by the MSCP (Ogden 1996). Specific objectives, as they relate to raptors, are as follows:

1. Document the protection of target species as specified in subarea plans and implementing agreements.
2. Document changes in preserved populations of covered species.
3. Describe new biological data collected.
4. Evaluate impacts of land uses and construction activities in and adjacent to the preserve.
5. Evaluate management activities and identify enforcement difficulties.

The purpose of this document is to provide guidance for consistency in the approach to surveying for raptors *during the breeding season and during the wintering period*. The below protocol is generic in nature but site-specific details, as to route, viewshed locations, and other important site features, are provided for each Raptor Monitoring Area (RMA) in Appendix C-1.

APPROACH

The following provides methodological details for the professional, with adequate raptor expertise, to conduct the breeding season and wintering period raptor monitoring in a consistent manner. The ability to detect trends (e.g., in raptor numbers, distribution, diversity, etc.) will be extremely important in order that adaptive management decisions be made in a timely manner. If trend analyses are to be interpretable, it is essential that the same locations within the preserve be monitored in a consistent manner. This would best be accomplished if the same individual or team monitored all locations, for all surveys.

ACRONYMS AND DEFINITIONS

Acronyms and definitions are attached (Appendix C-2). Use them consistently in order that there be continuity and clarity in all observations and record keeping.

SPECIES

Although all raptor species will be noted, there are eight MSCP, so-called “target,” raptor species: Bald Eagle (BE), Burrowing Owl (BO), Cooper’s Hawk (CH), Ferruginous Hawk (FH), Golden Eagle (GE), Northern Harrier (NH), American Peregrine Falcon (PF), and the Swainson’s Hawk (SH). Although you will not, necessarily, be searching for the BO at the most desirable time of day (early morning/early evening), any observations of BO or any other raptor species should be documented. Raptors will be the focus of the surveys but any observed sensitive species (regardless of taxa), interesting road kill, unusual biological observation, breeding colony, bird roost site, or other unique resource should also be noted on the WRI “Field Datasheet” (Appendix C-3).

TIMING AND FREQUENCY OF SURVEYS

Although it is common for ornithologists to identify a specific time of year as the “breeding season,” it is not possible to specify a timeframe, for our local raptors, that does not overlap with what is considered the wintering period. Because of the latitude of the MSCP, raptors are not restricted to a brief portion of the spring within which to breed. Many of our local raptors start breeding while other wintering and migrating raptors are still in the MSCP study area and environs. Therefore, the time of year that we call the “breeding season” could span December through August but varies considerably by species. Some GEs, for instance, can start nest building as early as December and still have nestlings in that nest as late as June. BOs, on the other hand, can start laying eggs in early April but fledge some young as late as August.

EQUIPMENT/SUPPLIES

Field vehicles should have 4WD capability if terrain requires. Binoculars, a camera, and a spotting scope of sufficient power for raptor observations are required. A magnification of 10X for binoculars and a range of approximately 20-60X for scopes are recommended. A cell phone may be very helpful in some locations, as could a set of “walkie-talkies” if more than one investigator will be in the field at the same time. Bring these survey guidelines, a copy of any authorization letters from resource agencies, any windshield placards (that indicates that you are under contract to conduct these surveys), local and project-generated site maps, and an adequate supply of “Transect Data Sheets” (Appendix C-3). To this, add your standard field equipment and supplies (field guides, hat, water, snacks, etc.). Although observers should be thoroughly familiar with all the local raptors, field guides that should be helpful include the Peterson guide, *Hawks* (Clark and Wheeler 1987) and the accompanying photographic guide (Wheeler and Clark 1999).

WEATHER

Monitoring should be conducted only during certain desirable weather conditions to maximize chances of documenting raptors. Inclement weather (rain, fog, winds greater than 20 mph, etc.) should be avoided. Occasional drizzle and winds up to 20 mph will not normally affect most raptor behavior. Observation in cold or wet weather should be done very carefully or completely discouraged. If an incubating bird is accidentally flushed during surveys, total nest failure could result for that season.

TIME OF DAY

The time of day, during which observations are made, is more important during the breeding season surveys than for the winter surveys (for most raptor species). Monitoring should take place from dawn through 1200, although professional experience may allow for some flexibility. Although BOs are not, necessarily, most active during this timeframe, you may note them and they should be documented as indicated below, as you would any raptor species. Since this is a crepuscular species, however, schedule sites that may support BOs for the early morning and/or early evening, whenever possible, to maximize chances of seeing this crepuscular species.

TWO TYPES OF OBSERVATIONS

Observations will be made two ways: (1) in vehicles, along established routes, and (2) at designated viewshed (i.e., observation) points. In addition, all reliable reports provided by interested individuals and cooperators will be verified and included in the data set but noted as “personal communications” with the appropriate documentation.

Vehicular Transects

Many of the breeding season raptor observations, and all those for the winter period, will be conducted from a vehicle. Therefore, vehicle speed will be an important variable. Speed will vary between road transects, depending on the road conditions, including traffic, and weather. That speed, however, should be consistent (year-to-year) for a particular transect in order that meaningful data comparisons can be made over time. Speed on a busy highway will have to be adequate to safely keep up with traffic. Some highway transects, that were deemed too dangerous, were removed from consideration. On a backcountry road, however, 10 mph may be the right speed. Safety should be the highest priority, and for that reason, an assistant to the driver is recommended to make observations and take notes, especially on busy roads.

Point/View shed Observations

Observation points have been established along some vehicle routes and at other desirable view shed locations for breeding season monitoring (see Appendix C-1). These will be especially important for riparian areas and inaccessible mountainous, and other, areas, where limited vehicle access prevents a reasonable survey of a RMA. At observation points along vehicle routes, a minimum of 10 minutes of actual observation is required. This means allowing whatever time is necessary to stop the vehicle in a safe, repeatable location, get out of the vehicle, and set up equipment (spotting scope, etc.) before starting the formal ten-minute observation (i.e., watching *and* listening). In situations where the observer is driving *through* the relevant habitat, a 5-minute observation period may be adequate. At some viewshed locations (like the top of a mountain), the observation time will be longer (perhaps 30 minutes). The most important issue here is that, once a viewing time period has been established for a particular RMA, it is maintained for consistency each year.

WHAT TO NOTE

All relevant data must be documented (see Transect Data Sheet, Appendix C-3). Sightings for *all* raptors will be documented. Note specific location of the raptor species *the first time it is observed* on each day of observation. Note age, sex, and any unusual plumage (if relevant) and

describe location(s) of any band(s) (metal right or metal left and sequence and numbers of any color bands), transmitter, or patagial wing markers. Avoid duplicate counts by noting unique characteristics of an individual and, when a bird is moving, its direction and relative speed. Record courtship and nesting behavior. If a nest is observed during the “winter” surveys, note its location on the topo map, what species of tree its in, height, size of nest, composition, and whether you consider it active. Indicate the basis for assumed activity (for instance, presence of an adult or pair near the nest, young, recent whitewash or greenery in /around nest).

CONTROL NUMBERING

Each control number for a study site and day of observation will be alphanumeric. For each species observed, the control number will start with the acronym for that species (see Appendix C-2) and be followed by “01.” The following control numbers, for that species, will end with 02, 03, etc., in the sequence in which the observations take place. This number is entered on the field data sheet (with all of its associated observations) and on the topo survey map, on which is always placed the survey date and the name(s) of the biologist(s). For instance, if the first observation of the day, at Mission Trails Regional Park, is a RT (Red-tailed Hawk), the control number will be “RT01.” The second RT will receive the control number “RT02.” If the next observation were a Cooper’s Hawk, it would be “CH01.” It will simplify records if each Transect Data Sheet and topo map is only used for one day’s observation at each site. However, there may be situations (such as when it takes more than one day to adequately survey a site or when it may lead to duplication or confusion later) when it makes sense to enter more than one day’s information on the same data sheet/map. It may also be beneficial to have all the breeding data on one map which keeps the picture in front of the observer at all times. This allows the observer to see gaps for certain species and explore areas not previously covered. The most important objective is to make sure the record is clear as to the date of each observation/set of observations and the name of the investigator so that clarification can be sought, if necessary.

Raptor, and other, nests are often less visible later in the breeding season, when deciduous trees have regained their foliage. However, note any stick nests in the area as “SN” followed by the appropriate observation number. Indicate on the data sheet if you know or suspect what species it belongs to and why. When summarizing yearly data, it will be important to determine which nests are alternate nests of the same pair and which represent additional pairs/territories. Do not get close enough to potentially disturb any nests, without approval from the Project Manager (PM) and Management Unit administrator.

Keep careful track of miles driven and times spent during vehicle transects and point location observations. Deduct any miles/time not spent on monitoring. These details are very important in order to allow data to be normalized over both time and distance to properly analyze for trends. There may be situations when you will not be able to track mileage or the miles you track are complicated by circling back through a study area to recheck a nest to confirm nesting, etc. Just keep good records that can be interpreted by someone else.

ENFORCEMENT/MANAGEMENT ISSUES

Note any enforcement or management problems or opportunities. Suggest corrective action or adaptive management, as appropriate, to the PM. Report any significant enforcement problems to the PM as soon as possible, but no later than within 24 hours of the observation.

RECORDS MANAGEMENT

Management of records is extremely important. Two-hole punched field forms and computer-generated project topo maps must be kept in Study Site folders (in a hard plastic or other secure file box provided) unless being copied. Field forms and topo maps must be attached to the inside of the Study Site folders using the two-hole clips at the end of each field day. Unless other provisions are made, field record copying should be done no less frequently than once a week, during the active field season, with copies placed in the appropriate administration project file for security.

THE SURVEYS

Breeding Season

In some management units, where a fulltime knowledgeable biologist is on staff, daily observations may be made, thereby providing greater potential for trend detection. However, the objective of these guidelines is to conduct up to 6 surveys at each of 12 RMAs (Figure C-1) for the breeding season raptor monitoring, where the assemblage of species dictates the actual number of replicates. Many stick nests will be located during the winter when the deciduous trees have lost their leaves. The next best opportunity to survey will often be early in the breeding season (December through April) when the adult raptors are establishing their territories and courting. Note that each species has a chronology for these behaviors. Some (like the GE, RT, and RS) will start breeding-related behaviors in December or January, while others (like the CH) may not display until April. At this time, they are obvious and concentrating their activities around the likely, and alternative, nest sites. In order to adequately characterize the raptor species present throughout the breeding season, the initial surveys at each site should be separated by 10-14 days, if possible. Subsequent surveys should be scheduled based on the raptor species present and where they are in their reproductive cycle. There will be a period, during which one of the adults will be incubating eggs or sheltering young, while the other adult is off hunting. During this time, it will be difficult to document many raptors and fieldwork may not be the best use of your time for that RMA. The next logical time to concentrate on conducting breeding season surveys will be when the young have fledged but are still dependent on the adults for food. At this time, there is a lot of activity and an increased chance of spotting a family unit because of the increased number of individuals per territory and, in some cases, the young will call attention to themselves by begging and/or calling to the parents.

The following times are recommended for the (breeding season) Raptor Monitoring Program:

- Late-December
- Mid-January
- Mid-February
- March
- Mid-April
- Mid-May

There are 12 RMAs that will be surveyed (Table C-1).

TABLE C-1. MSCP Raptor Monitoring Areas (Breeding Season)

<u>Area</u>	<u>Name</u>	<u>Study Sites* (original number(s))</u>
A	San Pasqual	San Pasqual (36), Lk. Hodges (7), Boden Cyn. (2), Rockwood (32)
B	Ramona Grasslands	Ramona Grasslands (30)
C	Penasquitos Canyon	Penasquitos Canyon (33)
D	Iron Mountain Complex	Iron Mountain** (11), San Vicente ((16), Route 67 (27)
E	San Diego River	San Diego River (26)
F	Sloan Canyon	Sloan Canyon (31), McGinty Mtn. North (5), Sycuan Mtn. North (17)
G	Sweetwater River	Sweetwater Reservoir (45), Rcho. S.D. East (42), Rcho. S.D. West (43), San Miguel Mtn. North (23)
H	Proctor Valley	Proctor Valley (25), San Miguel Mtn. South (23), Upper Otay Lk.(14)
I	Rancho Jamul	Jamul Ranch (3), Hollenbeck Canyon (34)
J	Border Fields	Border Fields (44), Tijuana River (part)
K	Brown Field Complex	Brown Field (22), Otay River, Spring Cyn. (part), Dennery Cyn. (part)
L	Otay Mountain	Otay Mountain (12), Marron Valley (13), Lower Otay Lake (14)

* In some cases, only a portion of a study site is included because of access, visibility, or some other reason (see detailed maps, Appendix C-1, for details).

** Including Monte Vista Ranch.

Each study site is followed by a number, which corresponds to the original study site number that was assigned to it (WRI 2002, 2004).

Winter Surveys

In keeping with the timing of many “winter” surveys (e.g., County Bird Atlas), the MSCP winter raptor surveys will occur primarily from *mid-December through February*, with possible changes in response to changes in weather conditions (i.e., global warming, cycles, etc.). This “winter” time period is somewhat arbitrary and we are not suggesting that raptors observed during this period are, necessarily, only birds that have migrated in and are wintering within the MSCP and environs. Similarly, the winter visit by some species may extend before and/or after this timeframe. The FH, for instance, can arrive on its MSCP wintering grounds by mid-September and not leave until mid-March. Many of the birds that you observe will be the same ones that you document during the “breeding season” surveys. The objective is to conduct three (3) vehicle-based surveys, along the coastal route depicted by Figure C-2. In order to adequately characterize the raptor species present throughout the winter season, the three surveys should be conducted according to the following schedule:

- Late December
- Mid-to-late January
- Mid-to-late February

Raptor, and other, nests are often more visible in the winter, when deciduous trees have lost their foliage. Knowledge about nest and breeding pair locations will help the monitor separate wintering birds from resident pairs. When summarizing yearly data, it will also be important to determine which nests are alternate nests of the same pair and which represent additional pairs/territories. Note any raptor nests in the area and/or if any nesting behavior is observed. Do not approach any nests, without approval from the PM and Management Unit administrator.

LITERATURE CITED

- Clark, W.S. and B.K. Wheeler. 1987. Peterson Field Guides—Hawks. Houghton Mifflin Company. Boston.
- County of San Diego. 1997. “Multiple Species Conservation Program”, County of San Diego; Subarea Plan. Adopted by the Board of Supervisors October 22, 1997.
- Ogden. 1996. “Biological Monitoring for the Multiple Species Conservation Program.” Prepared for the City of San Diego, California Department of Fish and Game, and U.S. Fish and Wildlife Service. Revised April 25.
- Wheeler, B.K and W.S. Clark 1999. A Photographic Guide to North American Raptors. Academic Press. San Diego.

APPENDIX C-2

ACRONYMS AND DEFINITIONS

Raptor and Corvid Species

AC	American crow
AK	American kestrel
BE*	BALD EAGLE
BH	Black hawk
BR	Barn owl
BO*	BURROWING OWL
CH*	COOPER'S HAWK
CR	Common raven
FH*	FERRUGINOUS HAWK
GE*	GOLDEN EAGLE
GO	Great-horned owl
HH	Harris' hawk
LO	Long-eared owl
MR	Merlin
NH*	NORTHERN HARRIER
OS	Osprey
PF*	PEREGRINE FALCON
PR	Prairie falcon
RS	Red-shouldered hawk
RT	Red-tailed hawk
SE	Short-eared owl
SO	Screech owl
SS	Sharp-shinned hawk
SH*	SWAINSON'S HAWK
TV	Turkey vulture
UA	Unidentifiable accipiter
UB	Unidentifiable buteo
UF	Unidentifiable falcon
UR	Unidentifiable raptor
WK	White-tailed kite
WH	White-tailed hawk
ZH	Zone-tailed hawk

Other Abbreviations

AB	Active burrow
Ad	Adult
CDFG	California Department of Fish and Game
CN	Cavity nest
F	Female
HY	Hatching year (when a bird is in its first year; i.e., the same calendar year as hatched).
Imm	Immature (a non-specific term that means "not adult").
M	Male
Mel	Melanistic (black/dark)
Ruf	Rufous/reddish
Sa	Sub adult (plumage that precedes adult plumage and appears much like it but with some characters that are not in adult plumage; used only for species, like the Golden Eagle, that can be distinguished at this age).
SN	Stick nest.
U	Unknown (e.g., unknown species, age, or sex).
USFWS	U.S. Fish and Wildlife Service

* MSCP target species.

APPENDIX C-3

TRANSECT DATA SHEET													
Wildlife Research Institute, Inc.							BIOLOGIST(S):						
		TIME (24hr)	Start	Finish	(minus time out) = TOTAL TIME:								
		TEMP (F):			OTHER WEATHER INFO.:								
DATE:	PAGE ___ OF ___	CLOUD CVR (%):			TRANSECT MILEAGE BEGIN:								
TRANSECT NAME & NUMBER:		WIND (mph):			TRANSECT MILEAGE END:								
		VISIBILITY (mi):			SUBTRACT MILEAGE:								
#		PRECIP:			TRANSECT TOTAL MILEAGE:								
WAYPOINTS (Start/End Points of Transects, Road Names, etc.)	SPECIES	TIME DURATION	SEX	AGE	PAIR	PERCHING	HUNTING	FEEDING	COURTSHIP	SOAPING	NESTING	COMMENTS, MILEAGE, TIME, ETC.	
1													
2													
3													
4													
5													
6													
7													
8													
9													
10													
11													
12													
13													
14													
15													
16													
17													
18													
19													
20													
21													
22													
23													
24													
25													
COMMENTS: (USE REVERSE FOR DRAWINGS OR ADDITIONAL NOTES)													